



ELSEVIER

Contents lists available at ScienceDirect

Data in Brief

journal homepage: www.elsevier.com/locate/dib

Data Article

Chemical and structural data of (1,2,3-triazol-4-yl)pyridine-containing coordination compounds

J. Conradie^{a,*}, M.M. Conradie^a, K.M. Tawfiq^{b,c}, M.J. Al-Jeboori^c, C. D'Silva^d, S.J. Coles^e, C. Wilson^f, J.H. Potgieter^{b,g}^a Department of Chemistry, University of the Free State, P.O. Box 339, Bloemfontein 9300, South Africa^b Division of Chemistry and Environmental Science, Manchester Metropolitan University, Manchester M1 5GD, UK^c Department of Chemistry, College of Education for Pure Science (Ibn Al-Haitham), University of Baghdad, Baghdad, Iraq^d Manipal University Jaipur, Department of Chemistry, VPO Dehmi Kalan, Jaipur 303007, Rajasthan, India^e EPSRC National Crystallography Service, School of Chemistry, University of Southampton, Southampton SO17 1BJ, England, UK^f School of Chemistry, University of Glasgow, Joseph Black Building, University Avenue, Glasgow G12 8QQ, Scotland, UK^g School of Chemical and Metallurgical Engineering, University of the Witwatersrand, Private Bag X3, Wits 2050, South Africa

ARTICLE INFO

Article history:

Received 31 March 2018

Accepted 24 August 2018

Available online 30 August 2018

ABSTRACT

The data presented in this paper are related to the research article entitled “Novel dichloro(bis{2-[1-(4-methylphenyl)-1H-1,2,3-triazol-4-yl-κN³]pyridine-κN})metal(II) coordination compounds of seven transition metals (Mn, Fe, Co, Ni, Cu, Zn and Cd)” (Conradie et al., 2018) [1]. This paper presents characterization and structural data of the 2-(1-(4-methyl-phenyl)-1H-1,2,3-triazol-1-yl)pyridine ligand (L²) (Tawfiq et al., 2014) [2] as well as seven dichloro(bis{2-[1-(4-methylphenyl)-1H-1,2,3-triazol-4-yl-κN³]pyridine-κN})metal(II) coordination compounds, [M(L²)₂Cl₂], all containing the same ligand but coordinated to different metal ions. The data illustrate the shift in IR, UV/VIS, and NMR (for diamagnetic complexes) peaks when L is coordinated to the metals, as well as the influence of the different metals on the peak positions. Solid state structural data is presented for M = Ni and Zn, while density functional theory calculated energies, structures and optimized coordinates

DOI of original article: <https://doi.org/10.1016/j.poly.2018.03.026>

* Corresponding author.

E-mail address: conradj@ufs.ac.za (J. Conradie).<https://doi.org/10.1016/j.dib.2018.08.125>2352-3409/© 2018 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

are provided for the lowest energy *cis* and *trans* conformations for L^2 as well as $[M(L^2)_2Cl_2]$ with $M = Mn, Fe, Co, Ni, Cu, Zn$ and Cd .

© 2018 The Authors. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

Specifications table

Subject area	Chemistry
More specific subject area	Coordination compounds
Type of data	Table, text file, graph, figure
How data was acquired	IR on Thermo-Nicolet FT-IR Spectrometer (AVATAR 320). Mass spectra on WATERS LCT premier mass spectrometer. Magnetic susceptibility with a Gouy magnetic susceptibility balance. X-ray structure on Rigaku SPIDER RAXIS image plate detector and Rigaku AFC12 goniometer equipped with an enhanced sensitivity (HG) Saturn724+ detector mounted at the window of an FR-E+ SuperBright molybdenum rotating anode generator with HF Varimax optics (100 μ m focus). NMR on an ECS-400 MHz, JEOL multi nuclear FT spectrometer. UV-vis spectra on a PerkinElmer Lambda 40 UV/Vis spectrometer. Electronic structure calculations using the Gaussian 09 package [3].
Data format	Raw, calculated, analyzed.
Experimental factors	–
Experimental features	–
Data source location	Division of Chemistry and Environmental Science, Manchester Metropolitan University, Manchester, M1 5GD, UK. Department of Chemistry, University of the Free State, Nelson Mandela Street, Bloemfontein, South Africa (DFT). Crystallographic data is held at the NCS University of Southampton. University of Sheffield (MS).
Data accessibility	Data is with article.
Related research article	J. Conradie, M.M. Conradie, K.M. Tawfiq, M.J. Al-Jeboori, S.J. Coles C. Wilson, J.H. Potgieter, Novel dichloro(bis[2-[1-(4-methylphenyl)-1H-1,2,3-triazol-4-yl- κ N ³]pyridine- κ N])metal(II) coordination compounds of seven transition metals (Mn, Fe, Co, Ni, Cu, Zn and Cd), Polyhedron, 2018, 151 (2018) 243-254. http://dx.doi.org/10.1016/j.poly.2018.03.026 .

Value of the data

- This data would be valuable for the further characterization and structural studies of (1,2,3-triazol-4-yl)pyridine-containing coordination compounds.
- This data provide NMR, IR, UV/VIS and magnetic moment data for (1,2,3-triazol-4-yl)pyridine-containing coordination compounds.
- MS fragmentation data for 2-(1-(4-methyl-phenyl)-1H-1,2,3-triazol-1-yl)pyridine ligand and seven dichloro(bis[2-[1-(4-methylphenyl)-1H-1,2,3-triazol-4-yl- κ N³]pyridine- κ N])metal(II) coordination compounds.
- This data provide solid state structures for two (1,2,3-triazol-4-yl)pyridine-containing coordination compounds.
- This data provide DFT optimized structures and coordinates for the lowest energy *cis* and *trans* isomers of the 2-(1-(4-methyl-phenyl)-1H-1,2,3-triazol-1-yl)pyridine ligand and seven (1,2,3-triazol-4-yl)pyridine-containing coordination compounds.

1. Data

1.1. Structural data

The $[M(L^2)_2Cl_2]$ compounds with $L^2 = 2-(1-(4\text{-methyl-phenyl})-1H-1,2,3\text{-triazol-1-yl})pyridine$, all have the same chemical formula $C_{28}H_{24}Cl_2N_8M$ with $M = Mn, Fe, Co, Ni, Cu, Zn$ and Cd . The X-ray solid state crystal structure of $[Ni(L^2)_2Cl_2]$ in Fig. 1 shows the coordination environment of the nickel metal ion with two 2-(1-(4-methyl-phenyl)-1H-1,2,3-triazol-1-yl)pyridine ligands L and two chlorides. A list of bond lengths and angles for the ligand L^2 (that crystallized together with $[Zn(L^2)_2Cl_2]$), $[Zn(L^2)_2Cl_2]$ and $[Ni(L^2)_2Cl_2]$ are listed in Table 1. The obtained geometrical parameters are in the




Fig. 1. View of $[Ni(L^2)_2Cl_2]$ showing the coordination environment Ni with two 2-(1-(4-methyl-phenyl)-1H-1,2,3-triazol-1-yl)pyridine ligands (L^2) and two chlorides, as well as the atom labelling scheme used in Table 1.

Table 1

Bond lengths [Å] and angles [°] for the ligand and compounds, obtained from solid state structural data [1]. Compound numbering according to Fig. 1.


$[Ni(L^2)_2Cl_2]$		$[Zn(L^2)_2Cl_2]$		L^2 (co-crystallize with $[Zn(L^2)_2Cl_2]$)	
Ni1–N1 ⁱ	2.1015(19)	Zn1–N1	2.144(3)		
Ni1–N1	2.1015(19)	Zn1–N1 ⁱ	2.144(3)		
Ni1–N8 ⁱ	2.0739(19)	Zn1–N8 ⁱ	2.191(4)		
Ni1–N8	2.0739(19)	Zn1–N8	2.191(4)		
Ni1–Cl1	2.4123(6)	Zn1–Cl1 ⁱ	2.4615(14)		
Ni1–Cl1 ⁱ	2.4123(6)	Zn1–Cl1	2.4615(14)		
N1–C2	1.341(3)	N1–C2	1.341(5)	N101–C102	1.270(16)
N1–C6	1.352(3)	N1–C6	1.346(5)	N101–C106	1.373(16)
N8–N9	1.316(3)	N8–N9	1.316(5)	N108–N109	1.315(11)
N8–C7	1.357(3)	N8–C7	1.363(5)	N108–C107	1.379(12)
N9–N10	1.352(3)	N9–N10	1.364(5)	N109–N110	1.378(12)
N10–C11	1.353(3)	N10–C11	1.352(5)	N110–C111	1.347(14)
N10–C12	1.428(3)	N10–C12	1.434(5)	N110–C112	1.433(13)
C2–C3	1.383(3)	C2–C3	1.385(6)	C102–C103	1.388(16)
C2–H2	0.9300	C2–H2	0.930	C102–H102	0.930
C3–C4	1.385(3)	C3–C4	1.381(6)	C103–C104	1.366(16)
C3–H3	0.9300	C3–H3	0.930	C103–H103	0.930
C4–C5	1.381(3)	C4–C5	1.378(6)	C104–C105	1.33(2)
C4–H4	0.9300	C4–H4	0.930	C104–H104	0.930
C5–C6	1.388(3)	C5–C6	1.403(6)	C105–C106	1.39(2)

Table 1 (continued)

[Ni(L ²) ₂ Cl ₂]		[Zn(L ²) ₂ Cl ₂]		L ² (co-crystallize with [Zn(L ²) ₂ Cl ₂])	
C5–H5	0.9300	C5–H5	0.930	C105–H105	0.930
C6–C7	1.460(3)	C6–C7	1.458(6)	C106–C107	1.451(14)
C7–C11	1.363(3)	C7–C11	1.369(6)	C107–C111	1.405(16)
C11–H11	0.9300	C11–H11	0.930	C111–H111	0.930
C12–C13	1.383(3)	C12–C13	1.376(6)	C112–C113	1.387(13)
C12–C17	1.384(3)	C12–C17	1.383(6)	C112–C117	1.338(16)
C13–C14	1.386(3)	C13–C14	1.397(6)	C113–C114	1.39(2)
C13–H13	0.9300	C13–H13	0.930	C113–H113	0.930
C14–C15	1.385(4)	C14–C15	1.385(7)	C114–C115	1.40(2)
C14–H14	0.9300	C14–H14	0.930	C114–H114	0.930
C15–C16	1.389(4)	C15–C16	1.388(7)	C115–C116	1.368(15)
C15–C18	1.508(3)	C15–C18	1.526(6)	C115–C118	1.502(14)
C16–C17	1.379(4)	C16–C17	1.390(7)	C116–C117	1.462(17)
C16–H16	0.9300	C16–H16	0.930	C116–H116	0.930
C17–H17	0.9300	C17–H17	0.930	C117–H117	0.930
C18–H18A	0.9600	C18–H18A	0.960	C118–H11A	0.960
C18–H18B	0.9600	C18–H18B	0.960	C118–H11B	0.960
C18–H18C	0.9600	C18–H18C	0.960	C118–H11C	0.960
N8i–Ni1–N8	180.0	N1–Zn1–Ni1	180.0		
N8–Ni1–N1i	100.41(8)	N1–Zn1–N8i	77.78(13)		
N8–Ni1–N1	79.59(8)	N1i–Zn1–N8	102.22(13)		
N1i–Ni1–N1	180.0	N8i–Zn1–N8	180.0		
N8 ⁱ –Ni1–Cl1	90.20(6)	N1–Zn1–Cl1 ⁱ	90.83(9)		
N8–Ni1–Cl1	89.80(6)	N1 ⁱ –Zn1–Cl1 ⁱ	89.17(9)		
N1 ⁱ –Ni1–Cl1	89.38(6)	N8 ⁱ –Zn1–Cl1 ⁱ	88.93(10)		
N1–Ni1–Cl1	90.62(6)	N8–Zn1–Cl1 ⁱ	91.07(10)		
N8 ⁱ –Ni1–Cl1 ⁱ	89.80(6)	N1–Zn1–Cl1	89.17(9)		
N8–Ni1–Cl1 ⁱ	90.20(6)	N1 ⁱ –Zn1–Cl1	90.83(9)		
N1 ⁱ –Ni1–Cl1 ⁱ	90.62(6)	N8 ⁱ –Zn1–Cl1	91.07(10)		
N1–Ni1–Cl1 ⁱ	89.38(6)	N8–Zn1–Cl1	88.93(10)		
Cl1–Ni1–Cl1 ⁱ	180.00(2)	Cl1 ⁱ –Zn1–Cl1	180.0		
C2–N1–Ni1	127.46(16)	C2–N1–Zn1	125.5(3)		
C6–N1–Ni1	114.55(15)	C6–N1–Zn1	115.4(3)		
N9–N8–Ni1	137.61(16)	N9–N8–Zn1	138.7(3)		
C7–N8–Ni1	112.60(15)	C7–N8–Zn1	111.1(3)		
C2–C3–C4	119.2(2)	C4–C3–C2	119.1(4)	C104–C103–C102	118.1(13)
C2–C3–H3	120.4	C4–C3–H3	120.4	C102–C103–H103	120.9
C2–N1–C6	117.9(2)	C2–N1–C6	119.0(4)	C102–N101–C106	119.4(12)
C3–C2–H2	118.8	C3–C2–H2	119.1	C103–C102–H102	118.5
C3–C4–H4	120.4	C3–C4–H4	120.0	C103–C104–H104	119.9
C4–C3–H3	120.4	C2–C3–H3	120.4	C104–C103–H103	120.9
C4–C5–C6	118.4(2)	C4–C5–C6	117.7(4)	C104–C105–C106	119.2(16)
C4–C5–H5	120.8	C4–C5–H5	121.1	C104–C105–H105	120.4
C5–C4–C3	119.2(2)	C5–C4–C3	120.1(4)	C105–C104–C103	120.2(14)
C5–C4–H4	120.4	C5–C4–H4	120.0	C105–C104–H104	119.9
C5–C6–C7	123.0(2)	C5–C6–C7	122.0(4)	C105–C106–C107	123.8(11)
C6–C5–H5	120.8	C6–C5–H5	121.1	C106–C105–H105	120.4
C7–C11–H11	127.7	C7–C11–H11	127.3	C107–C111–H111	127.4
C11–C7–C6	132.6(2)	C11–C7–C6	132.4(4)	C111–C107–C106	130.7(10)
C11–N10–C12	127.9(2)	C11–N10–C12	129.0(4)	C111–N110–C112	129.9(7)
C12–C13–C14	118.7(2)	C12–C13–C14	119.7(5)	C112–C113–C114	118.0(11)
C12–C13–H13	120.6	C12–C13–H13	120.2	C112–C113–H113	121.0
C12–C17–H17	120.3	C16–C17–H17	120.3	C112–C117–H117	120.9
C13–C12–C17	120.6(2)	C13–C12–C17	120.3(4)	C117–C112–C113	123.2(10)
C13–C12–N10	119.2(2)	C13–C12–N10	120.0(4)	C113–C112–N110	119.0(9)
C13–C14–H14	119	C13–C14–H14	119.5	C113–C114–H114	119.2
C14–C13–H13	120.6	C14–C13–H13	120.2	C114–C113–H113	121.0
C14–C15–C16	117.8(2)	C14–C15–C16	118.3(5)	C116–C115–C114	119.1(13)
C14–C15–C18	121.4(2)	C14–C15–C18	121.4(5)	C114–C115–C118	119.6(12)
C15–C14–C13	121.9(2)	C15–C14–C13	121.0(5)	C113–C114–C115	121.7(15)

Table 1 (continued)

[Ni(L ²) ₂ Cl ₂]		[Zn(L ²) ₂ Cl ₂]		L ² (co-crystallize with [Zn(L ²) ₂ Cl ₂])	
C15–C14–H14	119	C15–C14–H14	119.5	C115–C114–H114	119.2
C15–C16–H16	119.2	C17–C16–H16	119.3	C115–C116–H116	120.1
C16–C15–C18	120.8(2)	C16–C15–C18	120.4(5)	C116–C115–C118	121.2(10)
C16–C17–C12	119.4(2)	C12–C17–C16	119.5(5)	C112–C117–C116	118.3(12)
C16–C17–H17	120.3	C12–C17–H17	120.3	C116–C117–H117	120.9
C17–C12–N10	120.2(2)	C17–C12–N10	119.7(4)	C117–C112–N110	117.8(10)
C17–C16–C15	121.5(2)	C15–C16–C17	121.3(5)	C115–C116–C117	119.7(11)
C17–C16–H16	119.2	C15–C16–H16	119.3	C117–C116–H116	120.1
N10–C11–C7	104.5(2)	N10–C11–C7	105.3(4)	N110–C111–C107	105.2(8)
N10–C11–H11	127.7	N10–C11–H11	127.3	N110–C111–H111	127.4
N1–C2–C3	122.4(2)	N1–C2–C3	121.8(4)	N101–C102–C103	123.0(13)
N1–C2–H2	118.8	N1–C2–H2	119.1	N101–C102–H102	118.5
N1–C6–C5	122.9(2)	N1–C6–C5	122.2(4)	N101–C106–C105	120.0(12)
N1–C6–C7	114.2(2)	N1–C6–C7	115.8(4)	N101–C106–C107	116.2(10)
N8–C7–C11	108.4(2)	N8–C7–C11	107.7(4)	N108–C107–C111	107.8(9)
N8–C7–C6	119.1(2)	N8–C7–C6	119.8(4)	N108–C107–C106	121.5(8)
N8–N9–N10	106.02(18)	N8–N9–N10	106.1(3)	N108–N109–N110	107.9(8)
N9–N10–C11	111.3(2)	N9–N10–C11	110.7(3)	C111–N110–N109	110.2(7)
N9–N10–C12	120.81(19)	N9–N10–C12	120.2(3)	N109–N110–C112	119.9(9)
N9–N8–C7	109.79(19)	N9–N8–C7	110.2(4)	N109–N108–C107	108.9(8)
C15–C18–H18A	109.5	C15–C18–H18A	109.5		
C15–C18–H18B	109.5	C15–C18–H18B	109.5		
C15–C18–H18C	109.5	C15–C18–H18C	109.5		
H18A–C18–H18B	109.5	H18A–C18–H18B	109.5		
H18A–C18–H18C	109.5	H18A–C18–H18C	109.5		
H18B–C18–H18C	109.5	H18B–C18–H18C	109.5		
Symmetry transformations used to generate equivalent atoms	(i) -x+1,-y+1,-z+1	(i) -x,-y+1,-z+1		(i) -x,-y+1,-z+1	

**Fig. 2.** View of the free ligand L = 2-(1-(4-methyl-phenyl)-1H-1,2,3-triazol-1-yl)pyridine in the structure of [Zn(L²)₂Cl₂], L², disordered over an inversion centre, modelled as 0.5 occupied with isotropic displacement parameters.

same range as reported for related complexes [M(L¹)₂Cl₂] with L¹ = 2-(1-(4-methoxyphenyl)-1H-1,2,3-triazol-1-yl)pyridine and M = Co and Ni [4] and [Ni(L)₂Br₂] with L = 1-(cyclohexyl)-4-(2-pyridyl)-1,2,3-triazole [5] (Fig. 2). The obtained geometrical parameters for ligand L² (that crystallized together with [Zn(L²)₂Cl₂]), are in the same range as reported for ligand L², isolated alone [2].

1.2. Spectroscopic data

The UV/vis spectra of L² and the [M(L²)₂Cl₂] compounds are shown in Fig. 3 and characteristic data is summarized in Table 2. The IR spectra of L² and the [M(L²)₂Cl₂] compounds are shown in




Fig. 3. UV-vis spectra of L^2 and $[M(L^2)_2Cl_2]$ in DMSO solutions.

Table 2UV–vis spectral data and assignments of L^2 and $[M(L^2)_2Cl_2]$ in DMSO solutions.

Compound	Band Position λ_{\max} nm	Wave number (cm^{-1})	Extinction coefficient ϵ_{\max} ($dm^3 mol^{-1} cm^{-1}$)	Assignment
L^2	258, 287	38759, 34843	19740, 17200 (4×10^{-5} M)	Intra-ligand $\pi \rightarrow \pi^*$, $n \rightarrow \pi^*$
$[Mn(L^2)_2Cl_2]$	280, 284	35714, 35211	3165, 3124 (1×10^{-4} M)	Intra-ligand $\pi \rightarrow \pi^*$, $n \rightarrow \pi^*$
	682	14662	13	${}^6A_1g^{(S)} \rightarrow {}^4T_1g^{(4G)}$
$[Fe(L^2)_2Cl_2]$	284	35211	29513 (1.2×10^{-4} M)	Intra-ligand $\pi \rightarrow \pi^*$, $n \rightarrow \pi^*$
	326	30674	4947	CT
$[Co(L^2)_2Cl_2]$	280, 286, 298	35714, 34965, 33557	3672, 3347, 3240 (1×10^{-3} M)	Intra-ligand $\pi \rightarrow \pi^*$, $n \rightarrow \pi^*$
	615	1620	56	${}^4T_1g^{(F)} \rightarrow {}^4T_1g^{(P)}$
	677	14970	89	${}^4T_1g^{(F)} \rightarrow {}^4A_2g^{(F)}$
$[Ni(L^2)_2Cl_2]$	278, 282, 300	35971, 35460, 33333	3602, 3653, 3656 (1×10^{-3} M)	Intra-ligand $\pi \rightarrow \pi^*$, $n \rightarrow \pi^*$
	408	24509	20	${}^3A_2g^{(F)} \rightarrow {}^3T_1g^{(P)}$
	668	14970	8	${}^3A_2g^{(F)} \rightarrow {}^3T_1g^{(F)}$
$[Cu(L^2)_2Cl_2]$	279, 284	35842, 35211	3507, 3603 (1×10^{-3} M)	Intra-ligand $\pi \rightarrow \pi^*$, $n \rightarrow \pi^*$
	310	32258	3696	–
	908	11013	85	${}^2B_1g \rightarrow {}^2B_2g$
$[Zn(L^2)_2Cl_2]$	260, 287	38461, 35843	3220, 3067 (4×10^{-5} M)	Intra-ligand $\pi \rightarrow \pi^*$, $n \rightarrow \pi^*$
$[Cd(L^2)_2Cl_2]$	259, 287	38759, 34843	28005, 25695 (4×10^{-5} M)	Intra-ligand $\pi \rightarrow \pi^*$, $n \rightarrow \pi^*$

Fig. 4. Selected characteristic IR bands of L^2 and the $[M(L^2)_2Cl_2]$ compounds are listed and compared in reference [1]. The ionization data of the TOFMS-ES (+) mass spectra of L^2 and the $[M(L^2)_2Cl_2]$ compounds given are summarized in Table 3. The TOFMS-ES (+) mass spectra are provided in the Supplementary material. The 1H and ${}^{13}C$ NMR spectra of L^2 and the diamagnetic $[M(L^2)_2Cl_2]$ compounds ($M = Zn$ or Cd) are shown in Fig. 5, while data to determine the spin state (amount of unpaired d-electrons) for the paramagnetic $[M(L^2)_2Cl_2]$ compounds ($M = Mn, Fe, Co, Ni$ and Cu) are summarized in Table 4. More NMR spectra are provided in the Supplementary material.

1.3. DFT data

Both L^2 and the $[M(L^2)_2Cl_2]$ complexes may have different stereoisomers. The density functional theory calculated lowest energy *cis* and *trans* isomers, as well as the relative energies of the isomers, are shown in Fig. 6. The data associated with the geometry of the optimized geometries (Cartesian coordinates) of the compounds shown are provided in the Supplementary material.


Fig. 4. FT-IR absorption spectra of L² and [M(L²)₂Cl₂] compounds.

Table 3Fragmentation data of positive electrospray ionization of L^2 and the metals coordination compounds with ligand L^2 .

L^2 or coordination compound	Formula	MW	Fragmentation, m/z (%)
L^2	$C_{14}H_{12}N_4$	236.3	209 $[M-N_2]^+$ 22%, 237 $[M+H]^+$ 100%, 259 $[M+Na]^+$ 7%, 495 $[2M+Na]^+$ 18% (consist with literature [6])
$[Mn(L^2)_2Cl_2]$	$C_{28}H_{24}Cl_2MnN_8$	598.4	562.1 $[M-Cl]^+$ 90%, calculated for $[(C_{28}H_{24}N_8MnCl)]^+$, 237.1 $[L^2]^+$ 70%, calculated for $[(C_{14}H_{12}N_4)]^+$, 209.1 $[L^2-N_2]^+$ calculated for $[(C_{14}H_{14}N_2)]^+$ 100%
$[Fe(L^2)_2Cl_2]$	$C_{28}H_{24}Cl_2FeN_8$	599.3	563.1 $[M-Cl]^+$ 90%, calculated for $[(C_{28}H_{24}N_8FeCl)]^+$, 237.1 $[L^2]^+$ 70%, calculated for $[(C_{14}H_{12}N_4)]^+$, 209.1 $[L^2-N_2]^+$ calculated for $[(C_{14}H_{14}N_2)]^+$ 100%
$[Co(L^2)_2Cl_2]$	$C_{28}H_{24}Cl_2CoN_8$	602.4	566.1 $[M-Cl]^+$ 40%, calculated for $[(C_{28}H_{24}N_8CoCl)]^+$, 531 $[M-Cl_2]^+$ 5%, calculated for $(C_{28}H_{24}CoN_8)$
$[Ni(L^2)_2Cl_2]$	$C_{28}H_{24}Cl_2NiN_8$	602.1	565.1 $[M-Cl]^+$ 40%, calculated for $[(C_{28}H_{24}N_8NiCl)]^+$, 265 $[M-Cl_2-L^2+N_2]^+$ 50%, calculated for $[(C_{14}H_{10}N_2Ni)]^+$, 209.1 $[L^2-N_2]^+$ (10%), calculated for $[(C_{14}H_{12}N_2)]^+$
$[Cu(L^2)_2Cl_2]$	$C_{28}H_{24}Cl_2CuN_8$	607.0	594.1 $[M-Cl]^+$ 45%, calculated for $[(C_{28}H_{24}N_8CuCl)]^+$, 535.14 $[Cu(L^2)_2]^+$ 30%, calculated for $[C_{28}H_{24}CuN_8]^+$ 30%, 358 $[CuL^2]^+$, calculated for $[C_{14}H_{12}N_4CuCH_3COO^-]^+$ 100%, 237 $[L^2]^+$ calculated for $[(C_{14}H_{12}N_4)]^+$ 40%, 209 $[L^2-N_2]^+$ 15%, calculated for $[(C_{14}H_{12}N_2)]^+$
$[Zn(L^2)_2Cl_2]$	$C_{28}H_{24}Cl_2ZnN_8$	608.8	571.2 $[M-Cl]^+$ (80%), calculated for $[(C_{28}H_{24}N_8ZnCl)]^+$, 33 $[M-Cl-(L^2)]^+$ 5%, 237.1 $[L^2]^+$, calculated for $[(C_{14}H_{12}N_4Zn)]^+$ 30%, 209.1 $[L^2-N_2]^+$ calculated for $[(C_{14}H_{12}N_2)]^+$ 90%
$[Cd(L^2)_2Cl_2]$	$C_{28}H_{24}Cl_2CdN_8$	655.9	621.2 $[M-Cl]^+$ (100%), calculated for $[(C_{28}H_{24}N_8CdCl)]^+$, 237.1 $[L^2]^+$ (20%), calculated for $[(C_{14}H_{12}N_4)]^+$, 209.1 $[L^2-N_2]^+$ calculated for $[(C_{14}H_{12}N_2)]^+$ 50%

2. Experimental design, materials, and methods

Density functional theory (DFT) calculations were performed in the gas phase on the neutral compounds, using the B3LYP functional and the triple- ζ basis set 6–311 G(d,p) on all atoms except for Cd where the Stuttgart/Dresden (SDD) pseudopotential was used to describe the metal electronic core, while the metal valence electrons were described using the def2-TZVPP basis set [7]. The Gaussian 09 package [3] were used to optimize the compounds. The multiplicity used for L^2 and the $[M(L^2)_2Cl_2]$ compounds is singlet (L^2 , $[Zn(L^2)_2Cl_2]$ and $[Cd(L^2)_2Cl_2]$), doublet ($[Cu(L^2)_2Cl_2]$), triplet ($[Ni(L^2)_2Cl_2]$), quartet ($[Co(L^2)_2Cl_2]$), quintet ($[Fe(L^2)_2Cl_2]$) and sextet ($[Mn(L^2)_2Cl_2]$).




Fig. 5. ^1H and ^{13}C NMR spectrum of L^2 in CD_2Cl_2 , $[\text{Zn}(\text{L}^2)_2\text{Cl}_2]$ and $[\text{Cd}(\text{L}^2)_2\text{Cl}_2]$ in DMSO-d_6 .

Table 4

Data for determination of the spin state of paramagnetic $[\text{M}(\text{L}^2)_2\text{Cl}_2]$ complexes, μ_{eff} = effective magnetic moment.

Compound	amount of d electrons	μ_{eff} measured (B.M)	$\mu_{\text{eff}} = \sqrt{S(S+1)}$ calculated (B.M)	S
$[\text{Mn}(\text{L}^2)_2\text{Cl}_2]$	5	5.62	5.92	5/2
$[\text{Fe}(\text{L}^2)_2\text{Cl}_2]$	6	5.26	4.90	2
$[\text{Co}(\text{L}^2)_2\text{Cl}_2]$	7	3.98	3.87	3/2
$[\text{Ni}(\text{L}^2)_2\text{Cl}_2]$	8	3.00	2.83	1
$[\text{Cu}(\text{L}^2)_2\text{Cl}_2]$	9	1.70	1.73	1/2




Fig. 6. Density functional theory calculated optimized geometries of the lowest energy *cis* and *trans* isomers of L^2 and the $[M(L^2)_2Cl_2]$. The relative energies of the isomers, ΔE in eV, is also shown; the energy of the lowest energy isomer is indicated as 0 eV.

Acknowledgements

The National Mass Spectroscopy Centre at the University of Wales, Swansea is thanked for supplying the mass spectrometry data. XRD data and structures were supplied by the National Crystallography Service at the University of Southampton. KT expresses his gratitude to the Iraqi Government for financial support to conduct the research reported in the UK. This work has received support from the South African National Research Foundation (Grant numbers 113327 and 96111) and the Central Research Fund of the University of the Free State, Bloemfontein, South Africa. The High Performance Computing facility of the University of the Free State and the Centre for High Performance Computing CHPC of South Africa are gratefully acknowledged for computer time.

Transparency document. Supporting information

Transparency data associated with this article can be found in the online version at <https://doi.org/10.1016/j.dib.2018.08.125>.

Appendix A. Supporting information

Supplementary data associated with this article can be found in the online version at <https://dx.doi.org/10.1016/j.dib.2018.08.125>. CCDC 1813109 and 1813110 contains the supplementary crystallographic data for the crystals of this study. These data can be obtained free of charge via <http://www.ccdc.cam.ac.uk/conts/retrieving.html>, or from the Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: (+44) 1223-336-033; or e-mail: deposit@ccdc.cam.ac.uk.

References

- [1] J. Conradie, M.M. Conradie, K.M. Tawfiq, M.J. Al-Jeboori, S.J. Coles, C. Wilson, J.H. Potgieter, Novel Dichloro(bis(2-[1-(4-methylphenyl)-1H-1,2,3-triazol-4-yl- κ N²]pyridine- κ N})metal(II) coordination compounds of seven transition metals (Mn, Fe, Co, Ni, Cu, Zn and Cd), *Polyhedron* 151 (2018) 243–254. <https://doi.org/10.1016/j.poly.2018.03.026>.
- [2] K.M. Tawfiq, G.J. Miller, M.J. Al-Jeboori, P.S. Fennell, S.J. Coles, G.J. Tizzard, C. Wilson, J.H. Potgieter, Comparison of the structural motifs and packing arrangements of six novel derivatives and one polymorph of 2-(1-phenyl-1H-1,2,3-triazol-4-yl)pyridine, *Acta Cryst. B* 70 (2014) 379–389. <https://doi.org/10.1107/S2052520614001152>.
- [3] M.J. Frisch, G.W. Trucks, H.B. Schlegel, G.E. Scuseria, M.A. Robb, J.R. Cheeseman, G. Scalmani, V. Barone, B. Mennucci, G. A. Petersson, H. Nakatsuji, M. Caricato, X. Li, H.P. Hratchian, A.F. Izmaylov, J. Bloino, G. Zheng, J.L. Sonnenberg, M. Hada, M. Ehara, K. Toyota, R. Fukuda, J. Hasegawa, M. Ishida, T. Nakajima, Y. Honda, O. Kitao, H. Nakai, T. Vreven, J.A. Montgomery, Jr, J.E. Peralta, F. Ogliaro, M. Bearpark, J.J. Heyd, E. Brothers, K.N. Kudin, V.N. Staroverov, R. Kobayashi, J. Normand, K. Raghavachari, A. Rendell, J.C. Burant, S.S. Iyengar, J. Tomasi, M. Cossi, N. Rega, J.M. Millam, M. Klene, J.E. Knox, J.B. Cross, V. Bakken, C. Adamo, J. Jaramillo, R. Gomperts, R.E. Stratmann, O. Yazyev, A.J. Austin, R. Cammi, C. Pomelli, J.W. Ochterski, R. L. Martin, K. Morokuma, V.G. Zakrzewski, G.A. Voth, P. Salvador, J.J. Dannenberg, S. Dapprich, A.D. Daniels, Ö. Farkas, J. B. Foresman, J.V. Ortiz, J. Cioslowski, D.J. Fox, Gaussian 09, Revision D.01, Gaussian, Inc., Wallingford, CT, 2009.
- [4] J. Conradie, M.M. Conradie, K.M. Tawfiq, M.J. Al-Jeboori, S.J. Coles, C. Wilson, J.H. Potgieter, Synthesis, characterisation, experimental and electronic structure of novel Dichloro(bis(2-[1-(4-methoxyphenyl)-1H-1,2,3-triazol-4-yl- κ N³]pyridine- κ N))metal(II) compounds, metal = Mn, Co and Ni, *Journal of Molecular Structure*, 1161C: 89–99, DOI: [10.1016/j.molstruc.2018.02.036](https://doi.org/10.1016/j.molstruc.2018.02.036).
- [5] D. Schweinfurth, C.Y. Su, S.C. Wei, P. Braunsteind, B. Sarkar, Nickel complexes with “click”-derived pyridyl-triazole ligands: weak intermolecular interactions and catalytic ethylene oligomerisation, *Dalton Trans.* 41 (2012) 12984–12990. <https://doi.org/10.1039/C2DT31805A>.
- [6] F. Alonso, Y. Moglie, G. Radivoy, M. Yus, Click chemistry from organic halides, diazonium salts and anilines in water catalysed by copper nanoparticles on activated carbon, *Org. Biomol. Chem.* 9 (2011) 6385. <https://doi.org/10.1039/C1OB05735A>.
- [7] F. Weigend, R. Ahlrichs, Balanced basis sets of split valence, triple zeta valence and quadruple zeta valence quality for H to Rn: design and assessment of accuracy, *Phys. Chem. Chem. Phys.* 7 (2005) 3297–3305. <https://doi.org/10.1039/B508541A>.