

Carbon Dioxide Availability in Inlands Rivers Is Driven by Dissolved Organic Carbon, Not Warming: A Case Study of Tigris River

Batool Kadhim  , Mohammed Hamdan*  

Department of Biology, College of Science for Women, University of Baghdad, Baghdad, Iraq.

*Corresponding Author

Received 28/04/2023, Revised 01/07/2023, Accepted 03/07/2023, Published Online First 20/11/2023



This work is licensed under a [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/).

Abstract

Rivers became supersaturated with carbon dioxide (CO₂), so they play an essential role in the global carbon budgets. To increase our understanding of the source of CO₂ availability in rivers, we studied the role of climate-changed drivers of CO₂ availability, which are “dissolved organic carbon (DOC), and warming”. We sampled 45 locations of 3 parts within the Tigris River in Baghdad during autumn and winter. The results showed that all the studied variables (water temperature, pH, DOC, CO₂) changed over time. The variations in CO₂ availability are associated with changes in DOC concentration, not with water temperature. Overall, our results suggest that elevated CO₂ in rivers could result from increased DOC inputs. Therefore, we can conclude that increased DOC concentration in rivers was required for microbial respiration and photo-mineralization, which are the primary sources of CO₂ in river ecosystems.

Keywords: DOC, CO₂, Climate change, Warming, Carbon mineralization, Brownification.

Introduction

Dissolved organic matters are complicated and diverse organic components in nature that can pass through a 0.45 μm filter. They range from molecules with low molecular weight like; organic acid, amino acid, and sugar to high molecular weight molecules named hemic substances¹. Allochthonous organic matter (OM) is found in the soil of the surrounding terrestrial landscape as a result of the degradation of dead organisms² and as a product of plant photosynthesis, it can reach aquatic ecosystems due to ongoing climate change via increasing runoff due to high precipitation, on other hand autochthonous organic matter produced by aquatic algae³. Organic matters are different in their continent; for instance, organic matter formed from higher plants has a comparatively high proportion of aromatic carbon, a high phenolic content, and a low nitrogen concentration, while microbially derived organic matter (from bacteria and algae) has a lower

level of phenolics and an increased nitrogen concentration⁴⁻⁶.

Organic Matter is an essential source of carbon in aquatic ecosystems, e.g. dissolved organic carbon (DOC)⁷. High increasing inputs of dissolved organic matter (DOM) can cause a significant change in the physical and chemical properties of aquatic ecosystems like increasing carbon concentration concentrations⁸⁻¹⁰ via the brownification phenomenon.

Dissolved organic carbon has an essential role in regulating the carbon cycle¹¹, and is regarded as a significant contributor to increasing carbon concentration by bacterial and photo-mineralization of DOC to DIC (CO₂). Recently, aquatic ecosystems have received a high amount of CO₂ due to huge entrance of DOC¹². Rising DOC concentrations in water bodies lead to an increase in carbon continent¹³ and a decrease in oxygen level that causes hypoxia¹⁴.

¹⁵. More DOC concentration can lead to anoxia condition which in turn causes adverse effects on all aquatic life¹⁶. DOC effect on nutrient and light availability, with the gradient in concentration of DOC, high DOC concentration leads to increase associated nutrients that cause high biomass of algae which in turn causes shading and decreased light availability^{17, 18}.

Tigris River, a significant water source and one of Iraq's major rivers, it had been observed for a long time, and several studies had been conducted on it.

Material and Methods

Study Site Description

This study was conducted at some locations in the Tigris River in Baghdad (Fig. 1). The Tigris River, the second-largest river in western Asia, is surrounded by four nations: Iran, Iraq, Turkey, and Syria. The Tigris has a long history of the earliest known civilizations because of its significance to a primarily dry region and its value as a source of irrigation and transportation. Tigris River enters Baghdad city around 5 km from the tourist island in Baghdad; the town is split into two halves by the river, which runs from north to south: the right (Karkh) and the left (Risafa). Its breadth varies from 160 meters in straight sections to more than 400 meters in twisted branches, with a slope of 6.9 cm per kilometer. Numerous factors, including temperature variation and rainfall frequency, impact the Tigris River. Some of these factors are influenced by climatic change. Other human-caused variables can cause a wide range effecting on the Tigris River, e.g., irrigation, drainage work, and dams like the Samarra Dam, dams are significant to exploit the river as hydroelectric energy where they control the water level and maximize the flow potential of the river, in the same time dams cause increases in salinity concentration after the water return from these dams to the river. For our study, we sampled 45 locations in three parts on the Tigris River within Baghdad.

The estimated distance of the Tigris River between the sites is more than 10 km, where it is about 20-km from the first station to the third station, which are: Al-Gherai'at is a permanent island that is part of

Several researchers have examined changes in the chemical and physical characteristics monthly and seasonally and others studied alterations in nutrient concentrations and some investigated water quality of Tigris river¹⁹⁻²⁸. The analyses of carbon and its organic and inorganic sources are nonexistent in the Iraqi environment in general and on the Tigris River in particular. Based on a notable rise in carbon concentrations in aquatic ecosystems in last years²⁹, our study was concerned with studying organic carbon as a primary source of inorganic carbon, especially carbon dioxide in The Tigris River.

Rusafa and is northeast of Baghdad. It is distinguished for being covered with wild plants. Tigris River in this region occurs under direct sewage discharge from Al-Gherai'at's inhabited residential areas.

The second area is the Al-Shuhda'a Bridge, one of Baghdad's main bridges connecting the Karkh and Rusafa sides. Due to its high population and restaurants' spills of liquid pollutants into the river, it is classified as an active site.

The third area is Al-Jadriyah, a region on the side of Rusafa where the Tigris River makes a significant northeast turn. AL-Jadriyah has rapid urbanization and residential growth compared to the previously described sites, which produce a large volume and variety of liquid wastes.




Figure 1. Map of the studied sites (A-C) on the Tigris River in Baghdad. A) AL-Gherai'at (44°20' 11.2 E, 33°23' 11.1 N) B) AL-Shuhda'a Bridge (44°23' 10.0 E, 33°20' 18.9 N), and C) AL-Jadriyah (44°23' 07.4 E, 33°17' 26.5 N)

Field Measurements

Field measurements were performed using the standard methods in³⁰. Water temperature(°C) was measured by AL-Hana portable pocket thermometer. pH was determined with an HQ40 multimeter manufactured by HACH company. First, the device is calibrated using a buffer solution with pH values 9, 7, and 4. Then submerge the prop in water, the result is recorded once the reading is steady.

To estimate CO₂ availability, the method adopted by Hadi³¹⁾ and approved by Golterman³² was used for estimating CO₂ concentration in water. This method is based on the titration of 100 ml of water sample with a solution of sodium carbonate 0.2 N until pH equals 8.4. Then titration another 100 ml with Hydrochloric acid 0.1N until pH equals 4.2. After that, we calculate the total CO₂ in 100ml of the water sample by using this Eq.1:

$$X = ((A + B) * 4.4) \dots \dots \dots 1$$

Results and Discussion

Water temperature (mean ± SD) changed significantly during the study (Table 1) when it ranged between 19.7 ± 0.04 C in October and 7.3 ± 0.12 C in January (Fig. 2 a). pH (mean ± SD) also

Where A= Volume of titration with 0.2N Na₂CO₃, and B= Volume of titration with 0.1N HCl.

Then we calculated CO₂ by multiplying the value of X by 10, and the result was expressed by mg/L.

Laboratory Measurements

We collected the water sample and added H₂SO₄ until pH equaled 2.0 to remove DIC, then filtered the sample with a 0.45 Millipore filter in the field. The sample was transferred to the laboratory to calculate chemical oxygen demined (COD). In the lab, we used two solutions. The first was the digestion solution, which consists of 500 ml distilled water 10.216 ml of potassium dichromate, 167 ml of sulfuric acid, and 33.3 ml of mercuric sulfate. Then we completed the volume to liter. The second solution is an acid reagent consisting of 540 ml of H₂SO₄ and 5.5g of silver sulfate. After that we put the two solutions in the tube and added 2ml of samples to it. Then we put the tube in the digester for 120 minutes at 150 °C, then in the color meter, we read the result, and the result expressed as mg/l; later, we put the COD result in plutocalc water program then chose total organic carbon (TOC) which is the same represented DOC, because we filtered it firstly, as it is described in³³.

Statistics analyses

For statistical analyses (in SPSS 20) all variables were analyzed. T-tests estimated the differences variables between specific locations. A one-way ANOVA test was used to show if the variables differed significantly between the locations. Person's correlation coefficient (r) was used to show the correlation between water temperature and nutrient concentrations. Linear regression was used to indicate if the correlations were significant or not. We also found the mean and the standard divisions for all measured parameters.

changed during the study time (Table 1) when it ranged between 7.3 ± 0.16 in December and 8.4 ± 0.04 in January (Fig. 2d). Dissolved organic carbon (mean ± SD) changed significantly during the study

(Table 1) and ranged between 20.3 ± 9.3 mg/l, in January and 1.2 ± 1.2 mg/l in December (Fig. 2 b). Carbon dioxide concentration (mean \pm SD) changed significantly during the study (Table 1) and ranged between 204.1 ± 36.1 μ M in December and 110.2 ± 12.9 μ M in February (Fig. 2 c).

Carbon dioxide concentration was affected by dissolved organic carbon concentration during the study ($r=0.72$ and p -value <0.001) (Fig. 3 a) and not by water temperature during the study ($r=-0.24$ and p -value = 0.10) (Fig. 3 b).

Table 1. Two-way ANOVA analysis showing the effects of time, locations, and the interaction between them on the studied environmental variables.

Variable	Time		Location		Time x Location	
	F-value(df)	p-value	F-value(df)	p-value	F-value(df)	p-value
WT	3909.61(4, 44)	<0.0001	7.73(2,44)	0.001	2.55(8, 44)	0.02
pH	14.94(4, 44)	<0.0001	13.08(2, 44)	<0.0001	22.31(8, 44)	<0.0001
DOC	5.93(4, 44)	0.001	0.42(2, 44)	0.65	1.82(8, 44)	0.11
CO ₂	13.64(4, 44)	<0.0001	5.03(2, 44)	0.01	4.28(8, 44)	0.001




Figure 2. The studied environmental variables during the study time.




Figure 3. Carbon dioxide availability drivers during the study.

The results showed that water temperature, pH, DOC, and CO₂ concentration changed significantly during the study. It has been revealed statistically that CO₂ concentration was affected by DOC concentration and not by water temperature during the study. Carbon sources found in the aquatic environment either as dissolved organic carbon (DOC), which is created by photosynthesis and can come from the catchment adjacent to the water body or as dissolved inorganic carbon (DIC), especially CO₂, which is produced by the respiration of aquatic organisms and diffuses into water as atmospheric gas.

Carbon dioxide is an important substrate for an essential enzyme in the photosynthesis process^{34, 35}. Recently, CO₂ levels started to rise in the aquatic environment²⁹.

We noticed that DOC changed significantly during the study period. The high observed increase was in December and January because of high rainfall, which causes high surface runoff³⁶, as a result, carbon dioxide concentrations increased significantly due to the photolysis and biodegradation process of dissolved organic carbon^{37, 38}. On the other hand, the noticeable carbon dioxide levels during the study period were in AL-Gherai'at and AL-Shuhda'a bridge, and the reason is

due to anthropogenic activities such as the waste coming from the Baghdad Medical City hospital towards the AL-Shuhda'a bridge area, as well as different type of human activities like a discharge of sewage to the Tigris River without treatment near Al-Wazir Mosque, and the construction process of a bridge in AL-Gherai'at region that led to an increase in nutrients concentration e.g., phosphorus³⁹, which in turn causes the bacterial stimulation DOC-mineralization to CO₂⁴⁰.

Our results showed that the pH changes significantly following the change in carbon dioxide, where CO₂ is significant in controlling pH; pH falls as CO₂ rises and vice versa⁴¹⁻⁴³. Water temperature is an essential abiotic factor that drives a variety of characteristics of an organism's life, including respiration, fertility, growth, and nutrition, as well as its behavior and eventual survival. Heat plays an essential role in regulating respiration⁴⁴; when respiration processes noticeably increase due to rising temperatures, more CO₂ is produced. According to a different opinion, the heat had no apparent significant impact on metabolic activities⁴⁵, and the findings of our investigation agreed with this, where our results showed that temperature had no effect on CO₂ levels and the main drive of CO₂ concentrations was

dissolved organic carbon³⁶ that regulate the carbon budgets.

Recently, because of climate change, which has a significant role in increasing the concentration of

CO₂, it has been revealed that the increase in carbon dioxide is primarily related to increases in dissolved organic carbon concentration and not due to temperature variation.

Conclusion

Overall, our results revealed that warming does not influence carbon dioxide concentration clearly, whereas CO₂ availability is affected by allochthonous DOC inputs after the microbial and photo-mineralization processes of allochthonous and autochthonous dissolved organic carbon, which caused high levels of CO₂. We can conclude that climate change can control carbon dioxide concentration in aquatic ecosystems by increasing allochthonous DOC inputs from the lands

surrounding rivers due to high surface runoff processes brought on by precipitation due to the drought via increased global warming. This, in turn, raises the proportion of CO₂ availability in the river ecosystems. We recommend taking the different sources of carbon dioxide into account in the studies caring about human activities on the land surrounding the water bodies and their effects on the trophic status classification of aquatic ecosystems.

Acknowledgment

We thank the advanced aquatic lab staff in the department of Biology-College of Science for Women-University of Baghdad for providing the equipment we used. As well, the inorganic chemistry

lab staff in the Department of Chemistry-College of Science-University of Baghdad for providing access to the sample analysis in the laboratory.

Author's Declaration

- Conflicts of Interest: None.
- We hereby confirm that all the Figures and Tables in the manuscript are ours. Furthermore, any Figures and images, that are not ours, have been

- included with the necessary permission for re-publication, which is attached to the manuscript.
- Ethical Clearance: The project was approved by the local ethical committee in University of Baghdad.

Authors' Contributions Statement

The two authors designed the study; B.K. carried out the field and lab work; The data were analyzed statistically by M.H.; B.K wrote the first draft of the

manuscript.; M.H revised the final version of the manuscript.

References

1. Xiao K, Abbt-Braun G, Horn H. Changes in the characteristics of dissolved organic matter during sludge treatment: A critical review. *Water Res.* 2020; 187: 116441. <https://doi.org/10.1016/j.watres.2020.116441>.
2. Tiwari AK, Pal DB. Nutrients contamination and eutrophication in the river ecosystem. *Ecological Significance of River Ecosystems*: Elsevier Sci. 2022: 203-16. <https://doi.org/10.1016/B978-0-323-85045-2.00001-7>.
3. Wei M, Gao C, Zhou Y, Duan P, Li M. Variation in spectral characteristics of dissolved organic matter in inland rivers in various trophic states, and their relationship with phytoplankton. *Ecol Indic* 2019; 104: 321-32. <https://doi.org/10.1016/j.ecolind.2019.05.020>.

4. Liu S, Feng W, Song F, Li T, Guo W, Wang B, et al. Photodegradation of algae and macrophyte-derived dissolved organic matter: A multi-method assessment of DOM transformation. *Limnologia*. 2019; 77: 125683. <https://doi.org/10.1016/j.limno.2019.125683>.
5. Liu S, He Z, Tang Z, Liu L, Hou J, Li T, et al. Linking the molecular composition of autochthonous dissolved organic matter to source identification for freshwater lake ecosystems by combination of optical spectroscopy and FT-ICR-MS analysis. *Sci Total Environ*. 2020; 703: 134764. <https://doi.org/10.1016/j.scitotenv.2019.134764>.
6. Powers LC, Del Vecchio R, Blough NV, McDonald N, Schmitt-Kopplin P, Gonsior M. Optical properties and photochemical transformation of the dissolved organic matter released by Sargassum. *Front Mar Sci*. 2020;7:588287. <http://dx.doi.org/10.1029/2019GB006225>.
7. Wild B, Andersson A, Bröder L, Vonk J, Hugelius G, McClelland JW, et al. Rivers across the Siberian Arctic unearth the patterns of carbon release from thawing permafrost. *Proc Natl Acad Sci*. 2019; 116(21): 10280-5. <https://doi.org/10.1073/pnas.1811797116>.
8. Sterner RW, Reint KL, Lafrancois BM, Brovold S, Miller TR. A first assessment of cyanobacterial blooms in oligotrophic Lake Superior. *Limnol Oceanogr*. 2020; 65(12): 2984-98. <https://doi.org/10.1002/lno.11569>.
9. Kritzbeg ES, Hasselquist EM, Škerlep M, Löfgren S, Olsson O, Stadmark J, et al. Browning of freshwaters: Consequences to ecosystem services, underlying drivers, and potential mitigation measures. *Ambio*. 2020; 49: 375-90. <https://doi.org/10.1007/s13280-019-01227-5>.
10. Mackay E, Feuchtmayr H, De Ville M, Thackeray S, Callaghan N, Marshall M, et al. Dissolved organic nutrient uptake by riverine phytoplankton varies along a gradient of nutrient enrichment. *Sci Total Environ*. 2020; 722: 137837. <https://doi.org/10.1016/j.scitotenv.2020.137837>.
11. Liu F, Wang D, Zhang B, Huang J. Concentration and biodegradability of dissolved organic carbon derived from soils: A global perspective. *Sci Total Environ*. 2021; 754: 142378. <https://doi.org/10.1016/j.scitotenv.2020.142378>.
12. Hamdan M, Karlsson J, Byström P, Al-Haidarey MJ, Ask J. Carbon dioxide limitation of benthic primary production in a boreal lake. *Freshw Biol*. 2022; 67(10): 1752-60. <https://doi.org/10.1111/fwb.13972>
13. Chaplot V, Mutema M. Sources and main controls of dissolved organic and inorganic carbon in river basins: A worldwide meta-analysis. *J Hydrology*. 2021; 603: 126941. <https://doi.org/10.1016/j.jhydrol.2021.126941>.
14. Chen J, Li H, Zhang Z, He C, Shi Q, Jiao N, et al. DOC dynamics and bacterial community succession during long-term degradation of *Ulva prolifera* and their implications for the legacy effect of green tides on refractory DOC pool in seawater. *Water Res*. 2020; 185: 116268. <https://doi.org/10.1007/s11430-021-9954-0>
15. Ni W, Li M, Ross AC, Najjar RG. Large projected decline in dissolved oxygen in a eutrophic estuary due to climate change. *J Geophys Res: Oceans*. 2019; 124(11): 8271-89. <https://doi.org/10.1029/2019JC015274>.
16. Deeds J, Amirbahman A, Norton SA, Suito DG, Bacon LC. Predicting anoxia in low-nutrient temperate lakes. *Ecol Appl*. 2021; 31(6): e02361. <https://doi.org/10.1002/eap.2361>
17. Carrick HJ, Marble C, Tian YQ. Differential responses for stream algal assemblages exposed to factorial N and P enrichment along an in situ DOC gradient. *Front Environ Sci*. 2023; 11: 416. <https://doi.org/10.3389/fenvs.2023.1066586>.
18. Isles PD, Creed IF, Jonsson A, Bergström A-K. Trade-offs between light and nutrient availability across gradients of dissolved organic carbon lead to spatially and temporally variable responses of lake phytoplankton biomass to browning. *Ecosystems*. 2021: 1-16. <https://doi.org/10.1007/s10021-021-00619-7>
19. Al-Ani R, Al Obaidy A, Hassan F. Multivariate analysis for evaluation the water quality of Tigris River within Baghdad City in Iraq. *iraqi J Agric Sci*. 2019; 50(1): 331-42. <https://doi.org/10.36103/ijas.v50i1.299>
20. El-Amier YA, Al-Hadithy ON, Kadhim OG, El-Alfy M. Evaluation of Water and Sediment Quality of the Tigris River, Baghdad City, Iraq. *Am J Environ Sci* 2018; 1(1): 10-9. https://www.researchgate.net/publication/327476920_Evaluation_of_Water_and_Sediment_Quality_of_the_Tigris_River_Baghdad_City_Iraq.
21. Jabar SS, Hassan FM. Monitoring the Water Quality of Tigris River by Applied Overall Index of Pollution. *IOP Conf Ser : Environ Earth Sci*. 2022: IOP Publishing. <https://doi.org/10.1088/1755-1315/1088/1/012015>.
22. Majeed OS, Al-Azawi AJ, Nashaat MR. The Effect of Tharthar-Tigris Canal on the Environmental Properties of the Tigris River Northern Baghdad, Iraq. *Baghdad*

- Sci J. 2022; 19(6): 1177-1109.
<https://doi.org/10.21123/bsj.2022.6483>.
23. Mustafa MH, Jankeer MH. Quality Difference Between Two Location on Tigris River Within Mosul City. Rafidain J Sci. 2007; 1 (18) 111-124. (in Arabic).
<https://doi.org/10.33899/rjs.2007/166307>.
24. Saad MA, Antoine SE. Limnological studies on the River Tigris, Iraq. II. Seasonal variations of nutrients. Int Rev Hydrobiol. 1978; 63(5): 705-19.
<https://doi.org/10.1002/iroh.19780630516>.
25. Yasin S S. A Study of Monthly Changes in some Physical, Chemical, and Phytosanitary parameters in Tigris River at Salah Din Governorate SK Yasin. Tikrit J Agri Sci. 2018; 18(4): 131-141.
<https://www.iasj.net/iasj/download/06dc00574bba612c>.
26. Khadim HJ, Oleiwi HO, editors. Assessment of Water Quality in Tigris River of AL-Kut City, Iraq by Using GIS. E3S Web of Conferences; 2021; 2(1): 155-163: EDP Sciences.
<https://doi.org/10.1051/e3sconf/202131804001>.
27. Enad HY, Jaeel AJ, editors. Water quality index of Tigris River on Waist Governorate for aquatic life. IOP Conf Ser : Mater Sci Eng 2019: 12(3): 101-112. IOP Publishing. <https://doi.org/10.1088/1757-899X/584/1/012029>.
28. Al-Sudani IM. Water quality assessment of Tigris River using overall index of pollution (OIP). Baghdad Sci J. 2021; 18(2): 1099-100.
[https://doi.org/10.21123/B.S.J.2021.18.2\(Suppl.\).1095](https://doi.org/10.21123/B.S.J.2021.18.2(Suppl.).1095).
29. Weiss LC, Pötter L, Steiger A, Kruppert S, Frost U, Tollrian R. Rising pCO₂ in freshwater ecosystems has the potential to negatively affect predator-induced defenses in *Daphnia*. Curr Biol. 2018; 28(2): 327-32. e3. <https://doi.org/10.1016/j.cub.2017.12.022>.
30. APHA APHA. Standard Methods for the Examination of Water and Wastewater, 21st Ed Washington, DC.22621. pp .2005.
<https://doi.org/book/10.2105/SMWW.2882>.
31. Hadi RAM. Algae studies on the River Univ.College, Cardiff,UK; 1981.
[https://www.scirp.org/\(S\(351jmbntvnsjt1aadkposzje\)\)/reference/referencespapers.aspx?referenceid=533610](https://www.scirp.org/(S(351jmbntvnsjt1aadkposzje))/reference/referencespapers.aspx?referenceid=533610)
32. Golterman HL, Clymo R S, Ohnstad M A M. Methods for Physical and Chemical Analysis of Fresh Waters. 2nd ed. Oxford: Blackwell Scientific Publications; 1978.
<https://books.google.iq/books?id=JlgiAQAIAAJ>.
33. Williams CJ, Yamashita Y, Wilson HF, Jaffé R, Xenopoulos MA. Unraveling the role of land use and microbial activity in shaping dissolved organic matter characteristics in stream ecosystems. Limnol Oceanogr. 2010; 55(3): 1159-71.
<https://doi.org/10.4319/lo.2010.55.3.1159>.
34. Henard CA, Wu C, Xiong W, Henard JM, Davidheiser-Kroll B, Orata FD, et al. Ribulose-1, 5-bisphosphate carboxylase/oxygenase (RubisCO) is essential for growth of the methanotroph *Methylococcus capsulatus* strain Bath. Appl Environ. 2021; 87(18): e00881-21. <https://doi.org/10.1128/AEM.00881-21>.
35. Hassan FM, El-Sheekh MM, Wahhab TA. Environmental factors drive phytoplankton primary productivity in a shallow Lake. Egypt J Aquat Bio Fish. 2023; 27(2): 1-12.
<https://doi.org/10.21608/EJABF.2023.288620>.
36. Shatilla NJ, Carey SK. Assessing inter-annual and seasonal patterns of DOC and DOM quality across a complex alpine watershed underlain by discontinuous permafrost in Yukon, Canada. Hydrol Ear Syst Sci. 2019; 23(9): 3571-91. <https://doi.org/10.5194/hess-23-3571-2019>
37. Backlund P. Degradation of aquatic humic material by ultraviolet light. Chemosphere. 1992; 25(12): 1869-78.
38. Paine ER, Schmid M, Boyd PW, Diaz-Pulido G, Hurd CL. Rate and fate of dissolved organic carbon release by seaweeds: A missing link in the coastal ocean carbon cycle. J Phycol. 2021; 57(5): 1375-91.
[https://doi.org/10.1016/0045-6535\(92\)90026-N](https://doi.org/10.1016/0045-6535(92)90026-N).
39. Van Puijenbroek P, Beusen A, Bouwman A. Global nitrogen and phosphorus in urban waste water based on the Shared Socio-economic pathways. J Environ Manag. 2019; 231: 446-56.
<https://doi.org/10.1016/j.jenvman.2018.10.048>.
40. Allesson L, Andersen T, Dörsch P, Eiler A, Wei J, Hessen DO. Phosphorus availability promotes bacterial DOC-mineralization, but not cumulative CO₂-production. Front Microbiol. 2020; 11: 569879.
<https://doi.org/10.3389/fmicb.2020.569879>.
41. Liu J, Han G. Controlling factors of riverine CO₂ partial pressure and CO₂ outgassing in a large karst river under base flow condition. J Hydrol. 2021; 59(3): 125-132.
<https://doi.org/10.1016/j.jhydrol.2020.125638>
42. Sang L, Liu X, Sun D, Yang Y, Yang J, Wang Z, et al. Effect of tidal hydrology on soil anaerobic CO₂ production of freshwater marsh in the Yellow River estuary wetland, China. Ecol Indic. 2022; 145: 109747. <https://doi.org/10.1002/adma.201700990>.

43. Hamdan M, Byström P, Hotchkiss ER, Al-Haidarey MJ, Ask J, Karlsson J. Carbon dioxide stimulates lake primary production. *Sci Rep.* 2018; 8(1): 1-5. <https://www.nature.com/articles/s41598-018-29166-3>
44. Rasmusson LM, Buapet P, George R, Gullström M, Gunnarsson PC, Björk M. Effects of temperature and hypoxia on respiration, photorespiration, and photosynthesis of seagrass leaves from contrasting temperature regimes. *ICES J Mar Sci.* 2020; 77(6): 2056-65. <https://doi.org/10.1371/journal.pone.0083804>.
45. Huertas IE, Rouco M, Lopez-Rodas V, Costas E. Warming will affect phytoplankton differently: evidence through a mechanistic approach. *Proceedings of the Royal Society B: Biol sci.* 2011; 278(1724): 3534-43. <https://doi.org/10.1098/rspb.2011.0160>.

الكربون العضوي المذاب، ليس الاحترار، يقود توافر ثاني أكسيد الكربون في الأنهار الداخلية: حالة دراسية لنهر دجلة

بتول كاظم، محمد حمدان

قسم علوم الحياة، كلية العلوم للنبات، جامعة بغداد، بغداد، العراق.

الخلاصة

الانهار اصبحت مشبعة بثاني اوكسيد الكربون بشكل عالي وبذلك فهي تلعب دور مهم في كميات الكربون العالمية. لزيادة فهمنا حول مصادر الكربون المتوفرة في النظم البيئية النهرية، تم اجراء هذه الدراسة حول تأثير الكربون العضوي المذاب والحرارة (العوامل الرئيسية لتغير المناخ) كمحركات رئيسية لوفرة ثاني اوكسيد الكربون في الانهار. تم جمع العينات من خمسة واربعون موقع في ثلاثة اجزاء رئيسية لنهر دجلة داخل مدينة بغداد خلال فصلي الخريف والشتاء. اظهرت الدراسة ان جميع المتغيرات المدروسة (الحرارة، الأس الهيدروجيني، الكربون العضوي المذاب، وثاني اوكسيد الكربون) تتغير مع الوقت. كانت التغيرات في تركيز ثاني اوكسيد الكربون مرتبطة ايجابيا بالتغيرات في تركيز الكربون العضوي المذاب وليست بتغير درجات الحرارة. نتائجا بشكل عام تشير الى ان الزيادة بتراكيز ثاني اوكسيد الكربون في الانهار هو نتيجة لزيادة المدخلات من الكربون العضوي المذاب. وبذلك نستنتج ان الزيادة في تراكيز الكربون العضوي المذاب في الأنهار مطلوبة كمصدر لثاني اوكسيد الكربون من خلال عمليات التنفس الميكروبي والتحلل الكيميائي.

الكلمات المفتاحية: DOC، CO₂، تغير المناخ، الاحترار، تمعدن الكربون، تحول اللون البني.