We will study some methods that can be used to find the solutions of the first order equations which take the form y' = f(t, y)

<u>1- Separable Equations:</u>

Finding a way to separate the variables can be considered the best method to attempt first when trying to solve a differential equation. Even if we use one of the methods that we will discuss later for a given differential equation, we will invariably end up with the same integral to solve. Formally, a differential equation is separable if it can be written as:

$$\frac{dy}{dt} = a(t)b(y)$$

where $a, b: \mathbb{R} \to \mathbb{R}$ are continuous functions.

Solving Method:

The approach of solving a first order ODE using **separation of variables** is as follows:

Step 0: If required, we can first determine whether or not that the ODE is separable by using the theorem below:

<u>Theorem</u>: The DE y' = f(t, y) is separable if and only if $f(t, y) \frac{\partial^2 f}{\partial t \partial y} = \frac{\partial f}{\partial t} \frac{\partial f}{\partial y}$

Step 1: Rewrite (if necessary) the equation in the required form:

$$\frac{dy}{dt} = a(t)b(y)$$

Step 2: Find the general solution as follows:

$$\int \frac{dy}{b(y)} = \int a(t)dt$$

Step 3: If the initial conditions are given, solve to find the unique solution.

EXAMPLE: Verify that the IVP below is separable and find its solution:

$$y' = -2ty, y(0) = 1$$

Solution:

Step 0: To verify that the IVP above is separable, we set that y' = f(t, y) and we have to prove that: $f(t, y) \frac{\partial^2 f}{\partial t \partial y} = \frac{\partial f}{\partial t} \frac{\partial f}{\partial y}$

So
$$f(t, y) = -2ty$$
, $\frac{\partial f}{\partial t} = -2y$, $\frac{\partial f}{\partial y} = -2t$, and $\frac{\partial^2 f}{\partial t \partial y} = -2$
 $f(t, y) \frac{\partial^2 f}{\partial t \partial y} = -2ty$. $-2 = 4ty$ is equal to
 $\frac{\partial f}{\partial t} \frac{\partial f}{\partial y} = (-2y)$. $(-2t) = 4ty$

Step 1: $y' = -2ty \rightarrow \frac{dy}{dt} = -2ty \quad (a(t) = -2t, b(y) = y)$

Step2:
$$\rightarrow \frac{dy}{y} = -2tdt$$

 $\rightarrow \int \frac{dy}{y} = \int -2tdt \quad \rightarrow \quad Ln(y) = -t^2 + c \quad \rightarrow \quad e^{Ln(y)} = e^{-t^2 + c}$

The general solution is $y = e^{-t^2 + c}$

Step 3: $y = e^{-t^2+c}$ and y(0) = -1

So

 $y(0) = e^{-0^2 + c} = e^c = 1 \quad \to \ c = o$

Finally, $y = e^{-t^2+c} = e^{-t^2+0} = e^{-t^2}$ is the solution of the above IVP.

To verify the solution:

$$y = e^{-t^2} \rightarrow y' = e^{-t^2}(-2t) = -2ty \quad \text{correct} \ (\textcircled{O})$$

Homework: verify that each of the following DEs are separable and use separation of variables method to find the solution of each one:

1-
$$y' = 1 + y$$

2- $\frac{dy}{dt} = -\frac{y}{t}$, $y(0) = -4$
3- $\frac{1}{y^2}\frac{dy}{dx} + x^2 = 0$
4- $\frac{dy}{dt} = \frac{t(e^{t^2}+2)}{6y^2}$, $y(0) = 1$
5- $(y + y^2)dt - tdy = 0$, Hint: $\frac{rx+sy}{xy}$ can be transferred into $\frac{A}{x} + \frac{B}{y}$ for some
 $y(1) = 2$ real numbers A and B

EXAMPLE: Check if the ODE below is separable and find its solution:

$$(t-y)dt + tdy = 0$$

Solution: Step 0: First we need to rewrite the ODE in the form y' = f(t, y)

$$(t-y)dt + tdy = 0 \quad \rightarrow tdy = -(t-y)dt \quad \rightarrow tdy = (y-t)dt$$
$$\rightarrow \frac{dy}{dt} = \frac{y-t}{t} = \frac{y}{t} - 1 \quad \rightarrow \quad y' = \frac{y}{t} - 1 = f(t,y)$$

To determine if the ODE above is separable, we set that y' = f(t, y)and we have to prove that: $f(t, y) \frac{\partial^2 f}{\partial t \partial y} = \frac{\partial f}{\partial t} \frac{\partial f}{\partial y}$

So
$$f(t,y) = \frac{y}{t} - 1$$
, $\frac{\partial f}{\partial t} = -\frac{y}{t^2}$, $\frac{\partial f}{\partial y} = \frac{1}{t}$, and $\frac{\partial^2 f}{\partial t \partial y} = -\frac{1}{t^2}$

$$f(t, y) \frac{\partial^2 f}{\partial t \partial y} = \left(\frac{y}{t} - 1\right) \cdot -\frac{1}{t^2} = -\frac{y}{t^3} + \frac{1}{t^2} \text{ is NOT equal to } \frac{\partial f}{\partial t} \frac{\partial f}{\partial y} = -\frac{y}{t^2} \cdot \frac{1}{t} = -\frac{y}{t^3}$$

Therefore, the ODE is not separable.

<u>QUESTION</u>: Is it possible to overcome this problem, and make the above ODE is

separable?

ANSWER: YES, if it is Homogeneous Equation.

<u>2- Homogeneous Equations:</u>

An ordinary differential equation is said to be a homogeneous differential equation

if the following condition is satisfied

$$y' = f(zt, zy) = f(t, y) \quad \forall z \in \mathbb{R}$$

Let us come back to the previous example and see if it is homogenous

EXAMPLE: Check if the ODE is homogenous: (x - t)dt + tdy = 0 **Solution**: We need to show that y' = f(zt, zy) = f(t, y)We know that $y' = \frac{y}{t} - 1 = f(t, y)$ So, $f(zt, zy) = \frac{zy}{zt} - 1 = \frac{y}{t} - 1 = f(t, y)$

Therefore, the ODE is homogenous.

<u>QUESTION</u>: How can the method for solving homogeneous equations transform the not separable ODE into separable.

<u>ANSWER</u>: The method for solving homogeneous equations follows from this fact:

"The substitution y = vt (and therefore dy = vdt + tdv) transforms a homogeneous equation into a separable one."

Let us come back again to the previous example and see how is it going to be separable?

EXAMPLE: Transfer the not separable ODE below into separable by using the method of homogenous DE:

$$(t-y)dt + tdy = 0$$

Solution: we know that $y' = \frac{y}{t} - 1 = f(t, y)$

Set
$$y = vt \rightarrow y' = \frac{dy}{dt} = \frac{d(vt)}{dt} = v\frac{dt}{dt} + t\frac{dv}{dt} = v + tv' = f(t, vt)$$

 $\rightarrow tv' = f(t, vt) - v \rightarrow v' = \frac{f(t, vt) - v}{t}$
 $f(t, y) = \frac{y}{t} - 1 \rightarrow f(t, vt) = \frac{vt}{t} - 1 = v - 1$
 $\rightarrow v' = \frac{f(t, vt) - v}{t} = \frac{v - 1 - v}{t} = \frac{-1}{t}$

Now, this new form of our ODE $v' = \frac{-1}{t}$ is separable, so we can use the separation of variables method to solve it as follows:

$$v' = \frac{dv}{dt} = \frac{-1}{t} \quad \rightarrow \quad dv = -\frac{dt}{t} \quad \rightarrow \int dv = -\int \frac{dt}{t} \quad \rightarrow \quad v = -\ln(t) + c$$

But, $y = vt \rightarrow y = (-ln(t) + c)t$

To verify the solution:

$$y = (-\ln(t) + c)t \quad \to y' = (-\ln(t) + c) + t(-\frac{1}{t}) = (-\ln(t) + c) - 1 = \frac{y}{t} - 1$$

$$\rightarrow y' = \frac{y}{t} - 1$$

correct (☺♦)

<u>Strategy</u>: The approach of solving a first order DE using **the method of homogenous DE** is as follows:

Step 0: If required, check if the DE satisfies (**The homogeneous property**)

$$y' = f(zt, zy) = f(t, y)$$

Step 1: set that y = vt

Step 2: from f(t, y), find f(t, vt)

Step 3: substitute f(t, vt) in $v' = \frac{f(t, vt) - v}{t}$ to make the DE separable **Step 4:** use the separation of variable method to solve for v **Step 5:** substitute $v = \frac{y}{t}$ in the solution from step 4 to get the solution of the DE.

EXAMPLE: Verify that the ODE below is homogenous and find its solution:

 $(t^2 - y^2)dt + tydy = 0$

Step 0: To verify that the ODE above is separable, we set that y' = f(t,y) and we have to prove that y' = f(zt,zy) = f(t,y) $(t^2 - y^2)dt + tydy = 0 \rightarrow tydy = -(t^2 - y^2)dt \rightarrow \frac{dy}{dt} = \frac{y^2 - t^2}{ty}$ $f(t,y) = \frac{y^2 - t^2}{ty}$

$$f(zt, zy) = \frac{(zy)^2 - (zt)^2}{zt. zy} = \frac{z^2(y^2 - t^2)}{z^2 ty} = \frac{y^2 - t^2}{ty} = f(t, y)$$

The ODE is homogenous.

Step 1: set that y = vt

Step 2: from
$$f(t,y) = \frac{y^2 - t^2}{ty}$$
, we find $f(t,vt) = \frac{(vt)^2 - t^2}{t(vt)} = =$
$$\frac{t^2(v^2 - 1)}{t^2v} = \frac{v^2 - 1}{v}$$

Step 3: substitute $f(t, vt) = \frac{v^2 - 1}{v}$ in $v' = \frac{f(t, vt) - v}{t}$ to make the DE separable

$$v' = \frac{\frac{v^2 - 1}{v} - v}{t} = \frac{\frac{v^2 - 1 - v^2}{v}}{t} = \frac{-\frac{1}{v}}{t} = \frac{-1}{vt}$$

Step 4: use the separation of variable method to solve for v

$$v' = \frac{dv}{dt} = \frac{-1}{vt} \qquad \rightarrow v dv = -\frac{dt}{t} \rightarrow \int v dv = \int -\frac{dt}{t} \rightarrow \frac{v^2}{2} = -\ln(t) + c$$

$$\rightarrow v^2 = 2(-ln(t) + c)$$

Step 5: substitute $v = \frac{y}{t}$ in the solution from step 4 to get the solution of the ODE.

$$v^2 = 2(-ln(t) + c) \rightarrow \left(\frac{y}{t}\right)^2 = 2(-ln(t) + c) \rightarrow y^2$$

= $2t^2(-ln(t) + c)$

To verify the solution:

$$y^{2} = 2t^{2}(-\ln(t) + c) \quad \rightarrow 2yy' = 2t^{2}(\frac{-1}{t}) + (-\ln(t) + c)4t$$
$$\rightarrow y' = \frac{-2t + (-\ln(t) + c)4t}{2y} = \frac{2t(-1 + 2(-\ln(t) + c))}{2y} = \frac{t(-1 + \frac{y^{2}}{t^{2}})}{y} = \frac{t(\frac{-t^{2} + y^{2}}{t^{2}})}{y} = \frac{y^{2} - t^{2}}{ty}$$

1

correct (☺ы)

EXAMPLE: Verify that the ODE below is homogenous and find its solution:

$$(t^3 + y^3)dt - 2ty^2dy = 0$$

Step 0: To verify that the ODE above is separable, we set that y' = f(t,y) and we have to prove that y' = f(zt,zy) = f(t,y) $(t^3 + y^3)dt - 2ty^2dy = 0 \rightarrow 2ty^2dy = (t^3 + y^3)dt \rightarrow \frac{dy}{dt}$ $= \frac{(t^3 + y^3)}{2ty^2}$ $f(t,y) = \frac{(t^3 + y^3)}{2ty^2}$, $f(zt,zy) = \frac{((zt)^3 + (zy)^3)}{2zt(zy)^2} = \frac{z^3(t^3 + y^3)}{z^3(2ty^2)} = \frac{t^3 + y^3}{2ty^2} = f(t,y)$

The ODE is homogenous.

Step 1: set that y = vtStep 2: from $f(t, y) = \frac{t^3 + y^3}{2ty^2}$, we find $f(t, vt) = \frac{t^3 + (vt)^3}{2t(vt)^2} = \frac{t^3(1+v^3)}{t^3(2v^2)} = \frac{1+v^3}{2v^2}$ Step 3: substitute $f(t, vt) = \frac{1+v^3}{2v^2}$ in $v' = \frac{f(t,vt)-v}{t}$ to make the DE separable

$$v' = \frac{\frac{1+v^3}{2v^2} - v}{t} = \frac{\frac{1+v^3 - 2v^3}{2v^2}}{t} = \frac{\frac{1-v^3}{2v^2}}{t}$$

Step 4: use the separation of variable method to solve for v

$$v' = \frac{dv}{dt} = \frac{\frac{1-v^3}{2v^2}}{t}$$

$$\rightarrow \frac{dv}{\frac{1-v^3}{2v^2}} = \frac{dt}{t} \quad \rightarrow \quad \frac{2v^2dv}{1-v^3} = \frac{dt}{t} \quad \rightarrow \quad \int \frac{2v^2dv}{1-v^3} = \int \frac{dt}{t}$$

$$\rightarrow \quad \frac{-3}{-3} \int \frac{2v^2dv}{1-v^3} = \int \frac{dt}{t} \quad \rightarrow \quad \frac{2}{-3} \int \frac{-3v^2dv}{1-v^3} = \int \frac{dt}{t}$$

$$\rightarrow \quad \frac{2}{-3} \ln(1-v^3) = \ln(t) + c$$

Step 5: substitute $v = \frac{y}{t}$ in the solution from step 4 to get the solution of the ODE.

The solution of the ODE is $\frac{2}{-3}ln(1-(\frac{y}{t})^3) = ln(t) + c$ To verify the solution:

$$\frac{2}{-3}ln(1-(\frac{y}{t})^3) = ln(t) + c \rightarrow \frac{2}{-3}ln(1-\frac{y^3}{t^3}) = ln(t) + c$$

$$\rightarrow \frac{2}{-3}\frac{-(3\frac{y^2}{t^3}y'-3\frac{y^3}{t^4})}{1-\frac{y^3}{t^3}} = \frac{1}{t} \rightarrow \frac{2\frac{y^2}{t^3}y'-2\frac{y^3}{t^4}}{1-\frac{y^3}{t^3}} = \frac{1}{t} \rightarrow \frac{\frac{2ty^2y'-2y^3}{t^4}}{\frac{t^3-y^3}{t^3}} = \frac{1}{t}$$

$$\rightarrow \frac{2ty^2y'-2y^3}{t^3-y^3} = 1 \rightarrow 2ty^2y' - 2y^3 = t^3 - y^3 \rightarrow 2ty^2y' = t^3 + y^3$$

$$\rightarrow y' = \frac{(t^3 + y^3)}{2ty^2} \qquad \text{correct} (\textcircled{O} \textcircled{S})$$

Homework: verify that each of the following ODEs are homogenous and find the solution for each one:

EXAMPLE: Check if the ODE below is homogenous and find its solution:

$$2tydt + (t^2 - 1)dy = 0$$

SOLUTION:

Step 0: To verify that the ODE above is separable, we set that y' = f(t,y) and we have to prove that y' = f(zt,zy) = f(t,y) $2tydt + (t^2 - 1)dy = 0 \rightarrow (t^2 - 1)dy = -2tydt \rightarrow \frac{dy}{dt} = \frac{-2ty}{(t^2 - 1)}$ $f(t,y) = \frac{-2ty}{(t^2 - 1)}$ $f(zt,zy) = \frac{-2(zt)(zy)}{((zt)^2 - 1)} = \frac{-2z^2ty}{(z^2t^2 - 1)} \neq f(t,y)$

The ODE is nonhomogeneous.

<u>QUESTION</u>: Is it possible to overcome this problem, and make the above ODE is

solvable?

ANSWER: YES, if it is Exact Equation.

<u>3- Exact Equation:</u>

Consider the differential equation which takes the form

$$M(t, y)dt + N(t, y)dy = 0$$

we say that this differential equation is exact if it is satisfied this condition

$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial t}$$

Solving Method: The approach of solving a first order DE using **the method of Exact DE** is as follows:

Step 0: If required, check if the DE satisfies (The Exactness condition)

First, the equation should be in the form:

$$M(t, y)dt + N(t, y)dy = 0$$

and check it satisfies the condition: $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial t}$

Step 1: Assume that the function $\emptyset = \emptyset(t, y)$ (the solution of the general equation)

such that
$$\frac{\partial \phi}{\partial t} = M(t, y)$$
 and $\frac{\partial \phi}{\partial y} = N(t, y)$ (the old one)

(Which means that M(t, y)dt + N(t, y)dy = 0 becomes

$$\frac{\partial \emptyset}{\partial t} dt + \frac{\partial \emptyset}{\partial y} dy = 0$$

$$\rightarrow \quad \partial \emptyset = 0 \qquad \rightarrow \quad \emptyset = C \quad \forall t, y \in \mathbb{R}^2 \qquad)$$

Step 2: Integrate M(t, y) with respect of t to get:

$$\emptyset(t,y) = \int_t M(t,y)dt + h(y)$$

Step 3: Calculate the new $\frac{\partial \phi}{\partial y}$ as following:

$$\frac{\partial \phi}{\partial y} = \frac{\partial}{\partial y} \left(\int_t M(t, y) dt + h(y) \right) = \frac{\partial}{\partial y} \int_t M(t, y) dt + h'(y)$$

Step 4: Compare the new $\frac{\partial \phi}{\partial y}$ with the old $\frac{\partial \phi}{\partial y} = N(t, y)$ and solve to get h(y)

Step 5: Substitute h(y) in the equation from step 2:

$$\phi(t, y) = \int_t M(t, y)dt + h(y)$$

to get the solution of the DE.

Strategy: The steps of **the Exact DE method** are:

Step 0: Find M(t, y) and N(t, y) from M(t, y)dt + N(t, y)dy = 0

and verify that
$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial t}$$

Step 1: $A = \int M(t, y) dt$, with ignoring the integration constant, which should be a function of y.

Step 2: $B = \frac{\partial}{\partial v} A$

Step 3: $h(y) = \int (N(t, y) - B) dy$

Step 4: k = A + h(y) is the solution where k is constant.

EXAMPLE: Verify that the ODE is Exact and find its solution using the Exact

equation method:
$$2tydt + (t^2 - 1)dy = 0$$

Solution: Step 0: Find M(t, y) and N(t, y) from M(t, y)dt + N(t, y)dy = 0

and verify that
$$\frac{\partial M}{\partial y} = \frac{\partial N}{\partial t}$$

 $M(t, y) = 2ty$ and $N(t, y) = (t^2 - 1)$
 $\frac{\partial M}{\partial y} = 2t$ is equal to $\frac{\partial N}{\partial t} = 2t$ the ODE is Exact

Step 1: $A = \int M(t, y)dt$, $A = \int 2tydt = t^2y$, we integrate without the integration constant.

Step 2: $B = \frac{\partial}{\partial y}A$, $B = \frac{\partial}{\partial y}(t^2y) = t^2$ Step 3: $h(y) = \int (N(t, y) - B)dy$ $h(y) = \int (t^2 - 1 - t^2)dy = -\int dy = -y + c_1$ Step 4: k = A + h(y) is the solution where k is constant. $k = t^2y - y + c_1 \rightarrow t^2y - y = c$ where $c = k - c_1$ To verify the solution: $t^2y - y = c \rightarrow t^2y' + 2ty - yy' = 0 \rightarrow y' = \frac{dy}{dt} = \frac{2ty}{t^2 - 1}$ $\rightarrow 2tydt + (t^2 - 1)dy = 0$ correct (O O)

EXAMPLE: Verify that the ODE is Exact and find its solution using the Exact

equation method: $(e^{2y} - y\cos(ty))dt + (2te^{2y} - t\cos(ty) + 2y)dy = 0$

Solution:

Step 0: Find M(t,y) and N(t,y) from M(t,y)dt + N(t,y)dy = 0 and verify that $\frac{\partial M}{\partial y} = \frac{\partial N}{\partial t}$ $M(t,y) = e^{2y} - y \cos(ty)$ and $N(t,y) = 2te^{2y} - t \cos(ty) + 2y$ $\frac{\partial M}{\partial y} = 2e^{2y} - (y(-\sin(ty))t + \cos(ty)) = 2e^{2y} + ty \sin(ty) - \cos(ty))$ $\frac{\partial N}{\partial t} = 2e^{2y} - (t(-\sin(ty))y + \cos(ty)) = 2e^{2y} + ty \sin(ty) - \cos(ty))$

Step 1: $A = \int M(t, y) dt$,

Here, we find the integral with ignoring the integration constant.

$$A = \int (e^{2y} - y\cos(ty))dt = te^{2y} - \sin(ty)$$

Step 2:
$$B = \frac{\partial}{\partial y}A$$
,
 $B = \frac{\partial}{\partial y}(te^{2y} - sin(ty)) = 2te^{2y} - tcos(ty)$
Step 3: $h(y) = \int (N(t, y) - B)dy$
 $h(y) = \int (2te^{2y} - tcos(ty) + 2y - (2te^{2y} - tcos(ty)))dy$
 $= \int 2ydy = y^2 + c_1$

Step 4: k = A + h(y) is the solution where k is constant. $k = te^{2y} - sin(ty) + y^2 + c_1 \rightarrow te^{2y} - sin(ty) + y^2 = c$ where $c = k - c_1$

To verify the solution:

$$te^{2y} - sin(ty) + y^{2} = c$$

$$\rightarrow 2te^{2y}y' + e^{2y} - (tcos(ty)y' + ycos(ty) + 2yy' = 0)$$

$$\rightarrow y'(2te^{2y} - tcos(ty) + 2y) = -e^{2y} + ycos(ty)$$

$$\rightarrow y' = \frac{dy}{dt} = \frac{-e^{2y} + ycos(ty)}{(2te^{2y} - tcos(ty) + 2y)}$$

$$\rightarrow (e^{2y} - ycos(ty))dt + (2te^{2y} - tcos(ty) + 2y)dy = 0$$

$$correct (\textcircled{O})$$

<u>4- Solving the first-order Linear ODE by Integrating Factor method:</u>

We continue our quest for solutions of first-order differential equations by next examining linear equations.

<u>Def:</u> A first-order differential equation of the form

$$y' = a(t)y + b(t)$$

is said to be a linear equation in the variable y where $a: \mathbb{R} \to \mathbb{R}$ and $b: \mathbb{R} \to \mathbb{R}$.

Solving Method: The approach of solving a first order linear ODE using **the integrating factor** is as follows:

Step 0: Make sure that the equation is in the standard form:

$$y' = a(t)y + b(t)$$

Step 1: set $\mu = e^{-\int a(t)dt}$

This is called the **Integrating Factor**, and no constant need be used in evaluating the indefinite integral $\int a(t)dt$

Step 2: Multiply both sides of the equation in step 0 by μ , to get

$$\mu y' = \mu (a(t)y + b(t))$$

$$\rightarrow e^{-\int a(t)dt} \frac{dy}{dt} = e^{-\int a(t)dt} a(t)y + e^{-\int a(t)dt} b(t))$$

$$\rightarrow e^{-\int a(t)dt} \frac{dy}{dt} - e^{-\int a(t)dt} a(t)y = e^{-\int a(t)dt} b(t))$$

$$\rightarrow \frac{d}{dt} (e^{-\int a(t)dt} y) = e^{-\int a(t)dt} b(t)$$

$$\rightarrow d(e^{-\int a(t)dt} y) = e^{-\int a(t)dt}b(t)dt$$

Step 3: Integrate both sides of the equation above

$$\rightarrow \int d(e^{-\int a(t)dt} y) = \int e^{-\int a(t)dt} b(t)dt \rightarrow e^{-\int a(t)dt} y = \int e^{-\int a(t)dt} b(t)dt \rightarrow y = e^{\int a(t)dt} \int e^{-\int a(t)dt} b(t)dt$$

<u>Strategy Summary</u>: The steps of solving a first order linear DE using the integrating factor are:

Step 0: The equation should be in the form: y' = a(t)y + b(t) **Step 1:** set $\mu = e^{-\int a(t)dt}$ (integrate without constant) **Step 2:** The solution of the equation is

$$y = e^{\int a(t)dt} \int e^{-\int a(t)dt} b(t)dt$$

Step 3: Only for IVP, use the initial condition to find the value of the constant

<u>EXAMPLE</u>: Solve the differential equation $\frac{dy}{dt} - 3y = 0$

Solution:

Step 0: The equation should be in the form: y' = a(t)y + b(t) $\frac{dy}{dx} - 3y = 0 \rightarrow y' = 3y \rightarrow a(t) = 3, b(t) = 0$

Step 1: set $\mu = e^{-\int a(t)dt}$ (integrate without constant)

$$\mu = e^{-\int a(t)dt} = e^{-\int 3dt} = e^{-3t} \text{ the Integrating Factor}$$

Step 2: The solution of the equation is

$$y = e^{\int a(t)dt} \int e^{-\int a(t)dt} b(t)dt$$

$$y = e^{3t} \int e^{-3t} 0 dt = e^{3t} \int 0 dt = e^{3t} (0+c) = ce^{3t}$$
To varify the solution:

To verify the solution:

$$y = ce^{3t} \rightarrow \quad y' = 3ce^{3t} \rightarrow y' = 3y \qquad \text{ correct } (\textcircled{O} \textcircled{\diamond})$$

EXAMPLE: Solve the IVP
$$\frac{dy}{dt} - 2y = 6$$
, $y(0) = 9$

Solution:

Step 0: The equation should be in the form: y' = a(t)y + b(t) $\frac{dy}{dx} - 2y = 6 \rightarrow y' = 2y + 6 \rightarrow a(t) = 2, b(t) = 6$ Step 1: set $\mu = e^{-\int a(t)dt}$ (integrate without constant) $\mu = e^{-\int a(t)dt} = e^{-\int 2dt} = e^{-2t}$ the Integrating Factor Step 2: The solution of the equation is $y = e^{\int a(t)dt} \int e^{-\int a(t)dt} b(t)dt$ $y = e^{2t} \int e^{-2t} 6 dt = 6e^{2t} \int e^{-2t} dt = 6e^{2t} \left(\frac{e^{-2t}}{-2} + c\right) = -3 + 6ce^{2t}$

Step 3: Only for IVP, use the initial condition to find the value of the constant

$$y = -3 + 6ce^{2t} \rightarrow y(0) = -3 + 6c = 9 \rightarrow c = 2$$

$$\rightarrow y = -3 + 6ce^{2t} \quad \rightarrow \quad y = -3 + 12e^{2t}$$

To verify the solution:

$$y = -3 + 12e^{2t} \rightarrow y' = 24e^{2t} \rightarrow y' = 2(12e^{2t} - 3 + 3) = 2y + 6$$

correct (\textcircled{O}

EXAMPLE: Solve
$$t \frac{dy}{dt} - 4y = t^6 e^t$$

Solution:

Step 0: The equation should be in the form: y' = a(t)y + b(t)

$$t\frac{dy}{dt} - 4y = t^6 e^t \to y' = \frac{4}{t}y + t^5 e^t \to a(t) = \frac{4}{t}, b(t) = t^5 e^t$$

Step 1: set $\mu = e^{-\int a(t)dt}$ (integrate without constant)

$$\mu = e^{-\int a(t)dt} = e^{-\int \frac{4}{t}dt} = e^{-4ln(t)} = e^{ln(t^{-4})} = t^{-4}$$
 the Integrating Factor

Step 2: The solution of the equation is

$$y = e^{\int a(t)dt} \int e^{-\int a(t)dt} b(t)dt$$

 $y = t^4 \int t^{-4} t^5 e^t dt = t^4 \int t e^t dt$

We should use the integration by parts to solve
$$\int t e^t dt$$
 as follows:
Let $u = t$, $dv = e^t dt \rightarrow du = dt$, $v = \int e^t dt = e^t$
 $\rightarrow \int t e^t dt = uv - \int v du = te^t - \int e^t dt = te^t - e^t + c$
 $\rightarrow y = t^4 \int t e^t dt = t^4 (te^t - e^t + c)$

To verify the solution:

$$y = t^4(te^t - e^t + c) \rightarrow y' = t^4(te^t + e^t - e^t) + 4t^3(te^t - e^t + c)$$

$$\rightarrow y' = t^4(te^t) + 4t^3(te^t - e^t + c) = t^5e^t + \frac{4}{t}y$$

correct (☺ы́)

5- Nonlinear Equations can be transformed to linear by Substitutions:

We usually solve a differential equation by recognizing it as a certain kind of equation (say, separable, linear, or exact) and then carrying out a procedure, consisting of equation specific mathematical steps, that yields a solution of the equation. Sometimes a well-chosen substitution allows us to solve an equation. Sometimes it allows us to simplify an equation before we resort to numerical or qualitative techniques.

Next, we will study a type of differential equation, which take the general form:

$$g'(y)y' = a(t)g(y) + b(t)$$

Solving Method: The approach of solving this type of equations is as follows:

Step 0: Make sure that the equation is in the standard form

$$g'(y)y' = a(t)g(y) + b(t)$$

Step 1: Set v = g(y), which leads to $\frac{dv}{dt} = g'(y)y'$

Step 2: Substitute v, v' in the general equation, we get a linear equation with

respect to new dependent variable v

v' = a(t)v + b(t),

Step 3: Solve the last linear equation using integrating factor method to get v

Step 4: Use v = g(y), to get the solution in terms of y.

<u>EXAMPLE</u>: Solve $e^{y}y' + e^{y} = cos(t)$

Solution: The good choice is when we let

 $v = e^{y}$

$$v = e^{y} \rightarrow ln(v) = ln(e^{y}) \rightarrow ln(v) = y$$

$$\rightarrow v' = (e^{y})y'$$

So, $e^{y}y' + e^{y} = cos(t)$ becomes $v' + v = cos(t) \rightarrow v' = cos(t) - v$

Now, this equation is linear with respect to v, so we can solve it using

the integrating factor method.

<u>HW</u>: Finish the solution of the example above using the integrating factor method.

EXAMPLE: Solve $cos(y)y' = sin(y)t + 5e^{\frac{t^2}{2}}$ **Solution:** The good choice is when we let v = sin(y) $v = sin(y) \rightarrow v' = cos(y)y'$ and $y = sin^{-1}(v)$ $cos(y)y' = sin(y)t + 5e^{\frac{t^2}{2}}$ becomes $v' = vt + 5e^{\frac{t^2}{2}}$

Now, this equation is linear with respect to v, so we can solve it using

the integrating factor method.

<u>HW</u>: Finish the solution of the example above using the integrating factor method.

NOTE: Some differential equations may take different forms from the two examples above, So There are no general rules for finding good substitutions, see the following example:

<u>EXAMPLE</u>: Solve $y' = -e^y - 1$

Solution: The good choice is when we let $v = e^{-y}$

$$v = e^{-y} \rightarrow ln(v) = ln(e^{-y}) \rightarrow ln(v) = -y$$

$$\rightarrow y = -\ln(v) \qquad \rightarrow y' = -\frac{v'}{v}$$
$$y' = -e^y - 1 \quad \text{becomes} \quad -\frac{v'}{v} = -\frac{1}{v} - 1 \quad \rightarrow v' = 1 + v$$

Now, this equation is linear with respect to v, so we can solve it using

the integrating factor method.

Step 0: The equation should be in the form: y' = a(t)y + b(t) $v' = 1 + v \longrightarrow a(t) = 1, b(t) = 1$ **Step 1:** set $\mu = e^{-\int a(t)dt}$ (integrate without constant) $u = e^{-\int 1dt} = e^{-\int dt} = e^{-t}$ the Integrating Factor Step solution equation 2: The of the is $v = e^{\int a(t)dt} \int e^{-\int a(t)dt} b(t)dt$ $v = e^t \int e^{-t}(1) dt = e^t \int e^{-t} dt = e^t (-e^{-t} + c) = -1 + ce^t$ $v = -1 + ce^t$ but $v = e^{-y}$ $v = -1 + ce^t \rightarrow e^{-y} = -1 + ce^t$ $\rightarrow \ln(e^{-y}) = ln(-1+ce^t) \quad \rightarrow -y = ln(-1+ce^t)$ $\rightarrow v = -ln(-1+ce^{t})$

To verify the solution:

$$y = -ln(-1+ce^{t}) \rightarrow y' = \frac{-ce^{t}}{(-1+ce^{t})} = \frac{-(ce^{t}-1)-1}{e^{-y}} = \frac{-e^{-y}-1}{e^{-y}} = \frac{-e^{-y}}{e^{-y}} - \frac{1}{e^{-y}}$$
$$\rightarrow y' = -1 - e^{y} \qquad \text{correct} (\textcircled{O})$$

6-Bernoulli's Equation:

<u>Def</u>: The differential equation $y' = a(t)y + y^nb(t)$

where n is any real number ($n \neq 0$ and $n \neq 1$), is called Bernoulli's equation.

<u>Note</u>: For n = 0 and n = 1, the equation above is linear.

<u>Strategy:</u> To solve Bernoulli's equation:

- 1- Do the substitution $z = y^{1-n}$ because this substitution reduces any equation of the form above to a linear equation with respect to z.
- 2- Solve the linear equation in terms of z to find z using the integrating factor method.
- 3- Use $z = y^{1-n}$ to make the solution in terms of y.
- **<u>EXAMPLE</u>**: Solve the IVP $t \frac{dy}{dt} + y = t^2 y^2$, y(1) = 1

Solution: $t\frac{dy}{dt} + y = t^2y^2 \rightarrow ty' + y = t^2y^2 \rightarrow y' = -\frac{1}{t}y + ty^2$

This equation is Bernoulli's equation with n=2

Let $z = y^{1-n} = y^{1-2} = y^{-1} \rightarrow y = z^{-1} \rightarrow y' = -z^{-2}z'$ $y' = -\frac{1}{t}y + ty^2$ becomes $-z^{-2}z' = -\frac{1}{t}z^{-1} + t(z^{-1})^2$ $\rightarrow -z^2(-z^{-2}z') = -z^2(-\frac{1}{t}z^{-1} + tz^{-2})$ $\rightarrow z' = \frac{1}{t}z - t$

Now, this equation is linear with respect to z, so we can solve it using **the integrating factor method.**

Step 0: The equation should be in the form: y' = a(t)y + b(t)

$$z' = \frac{1}{t}z - t \quad \rightarrow a(t) = \frac{1}{t}, b(t) = -t$$

Step 1: set $\mu = e^{-\int a(t)dt}$ (integrate without constant)
 $\mu = e^{-\int \frac{1}{t}dt} = e^{-ln(t)} = e^{ln(t^{-1})} = t^{-1}$ the Integrating Factor
Step 2: The solution of the equation is
 $z = e^{\int a(t)dt} \int e^{-\int a(t)dt} b(t)dt$
 $z = t \int t^{-1}(-t) dt = t \int -dt = t(-t+c) = -t^2 + ct$
 $z = -t^2 + ct$ but $y = z^{-1} \rightarrow z = y^{-1}$

$$z = -t^2 + ct \rightarrow y^{-1} = -t^2 + ct \rightarrow y = \frac{1}{-t^2 + ct}$$

Step 3: Only for IVP, use the initial condition to find the value of the constant

$$y = \frac{1}{-t^2 + ct} \rightarrow y(1) = \frac{1}{-1^2 + c(1)} = 1 \rightarrow c = 2$$

 $\rightarrow y = \frac{1}{-t^2 + 2t}$

To verify the solution:

$$y = \frac{1}{-t^2 + 2t} \rightarrow \quad y' = \frac{-(-2t+2)}{(t^2 + 2t)^2} = \frac{-\frac{1}{t}(-t^2 + 2t) + t}{\frac{1}{y^2}} = \frac{-\frac{1}{t}\frac{1}{y} + t}{\frac{1}{y^2}}$$
$$\rightarrow \quad y' = \left(-\frac{1}{t}\frac{1}{y} + t\right)y^2 = -\frac{1}{t}y + ty^2 \qquad \text{correct} \ (\textcircled{O})$$

EXAMPLE: Solve
$$\frac{dy}{dt} - \frac{y}{3t} = e^t y^4$$

Solution: $\frac{dy}{dt} - \frac{y}{3t} = e^t y^4 \rightarrow y' = \frac{y}{3t} + e^t y^4$

This equation is Bernoulli's equation with n=4

Let
$$z = y^{1-n} = y^{1-4} = y^{-3} \rightarrow y = z^{-\frac{1}{3}} \rightarrow y' = -\frac{1}{3}z^{-\frac{4}{3}}z'$$

 $y' = \frac{y}{3t} + e^t y^4$ becomes $-\frac{1}{3}z^{-\frac{4}{3}}z' = \frac{z^{-\frac{1}{3}}}{3t} + e^t(z^{-\frac{1}{3}})^4$
 $\rightarrow -3z^{\frac{4}{3}}(-\frac{1}{3}z^{-\frac{4}{3}}z') = -3z^{\frac{4}{3}}(\frac{z^{-\frac{1}{3}}}{3t} + e^tz^{-\frac{4}{3}})$
 $\rightarrow z' = -\frac{1}{t}z - 3e^t$

Now, this equation is linear with respect to z, so we can solve it using **the integrating factor method.**

Step 0: The equation should be in the form: y' = a(t)y + b(t) $z' = -\frac{1}{t}z - 3e^t \rightarrow a(t) = -\frac{1}{t}, b(t) = -3e^t$

Step 1: set $\mu = e^{-\int a(t)dt}$ (integrate without constant)

$$\mu = e^{-\int -\frac{1}{t}dt} = e^{\ln(t)} = t$$
 the Integrating Factor

Step 2: The solution of the equation is

$$z = e^{\int a(t)dt} \int e^{-\int a(t)dt} b(t)dt$$

 $z = t^{-1} \int t(-3e^t) dt = -3t^{-1} \int te^t dt$

We should use the integration by parts to solve $\int -3t e^t dt$ as follows:

Let
$$u = t$$
, $dv = e^t dt \rightarrow du = dt$, $v = \int e^t dt = e^t$
 $\rightarrow \int t e^t dt = uv - \int v du = te^t + \int e^t dt = te^t - e^t + c$
 $\rightarrow z = -3t^{-1} \int te^t dt = -3t^{-1}(te^t - e^t + c)$
 $z = -3t^{-1}(te^t - e^t + c)$ but $z = y^{-3}$
 $\rightarrow y^{-3} = -3t^{-1}(te^t - e^t + c)$
 $\rightarrow y^3 = \frac{-t}{3(te^t - e^t + c)}$

To verify the solution:

$$y^{3} = \frac{-t}{3(te^{t} - e^{t} + c)} \rightarrow 3y^{2}y' = \frac{3(te^{t} - e^{t} + c) + t(3(te^{t} + e^{t} - e^{t}))}{(3(te^{t} - e^{t} + c))^{2}} = \frac{-3(te^{t} - e^{t} + c) + 3t^{2}e^{t}}{(3(te^{t} - e^{t} + c))^{2}}$$

$$\rightarrow 3y^{2}y' = \frac{\frac{t}{y^{3}} + 3t^{2}e^{t}}{(\frac{-t}{y^{3}})^{2}} = \frac{3\frac{t^{2}}{y^{3}}(\frac{1}{3t} + e^{t}y^{3})}{\frac{t^{2}}{y^{6}}} = \frac{3(\frac{1}{3t} + e^{t}y^{3})}{\frac{1}{y^{3}}} = 3y^{3}(\frac{1}{3t} + e^{t}y^{3})$$

$$\rightarrow y' = \frac{y}{3t} + e^{t}y^{4} \qquad \text{correct} (\textcircled{S})$$

EXAMPLE: Solve $y' + \frac{1}{10}y - \cos(t)y^2 = 0$

Solution: $y' + \frac{1}{10}y - \cos(t)y^2 = 0 \quad \rightarrow \quad y' = -\frac{1}{10}y + \frac{1}{10}y + \frac$

 $cos(t)y^2$

This equation is Bernoulli's equation with n=2

Let $z = y^{1-n} = y^{1-2} = y^{-1} \to y = z^{-1} \to y' = -z^{-2}z'$

$$y' = -\frac{1}{10}y + \cos(t)y^{2} \text{ becomes}$$

$$-z^{-2}z' = -\frac{1}{10}z^{-1} + \cos(t)(z^{-1})^{2}$$

$$\rightarrow -z^{2}(-z^{-2}z') = -z^{2}(-\frac{1}{10}z^{-1} + \cos(t)z^{-2})$$

$$\rightarrow z' = \frac{1}{10}z - \cos(t)$$

Now, this equation is linear with respect to z, so we can solve it using **the integrating factor method.**

Step 0: The equation should be in the form: y' = a(t)y + b(t)

$$z' = \frac{1}{10}z - cos(t) \rightarrow a(t) = \frac{1}{10}, b(t) = -cos(t)$$

Step 1: set $\mu = e^{-\int a(t)dt}$ (integrate without constant)

$$\mu = e^{-\int \frac{1}{10}dt} = e^{-\frac{t}{10}}$$
 the Integrating Factor

Step 2: The solution of the equation is $z = e^{\int a(t)dt} \int e^{-\int a(t)dt} b(t)dt$ $z = e^{\frac{t}{10}} \int e^{-\frac{t}{10}} (\cos(t)) dt$

We should use the integration by parts twice to solve $\int e^{-\frac{t}{10}} cos(t) dt$ as follows:

First, we let u = cos(t), $dv = e^{-\frac{t}{10}} dt$

$$\rightarrow du = -\sin(t)dt$$
, $v = \int e^{-\frac{t}{10}} dt = -10e^{-\frac{t}{10}}$

$$\rightarrow \int e^{-\frac{t}{10}} \cos(t) \, dt = uv - \int v \, du = -10\cos(t)e^{-\frac{t}{10}} - \int -10e^{-\frac{t}{10}}(-\sin(t))dt$$
We should use the integration by parts again to solve $\int -10e^{-\frac{t}{10}}(-\sin(t))dt$

Now, we let u = -sin(t), $dv = -10e^{-\frac{t}{10}} dt$

 $100\int e^{-\frac{t}{10}}\cos(t)\,dt)$

$$= -10\cos(t)e^{-\frac{t}{10}} + 100\sin(t)e^{-\frac{t}{10}} -$$

$$100 \int e^{-\frac{t}{10}} \cos(t) dt)$$

$$\rightarrow 101 \int e^{-\frac{t}{10}} \cos(t) dt = -10 \cos(t) e^{-\frac{t}{10}} + 100 \sin(t) e^{-\frac{t}{10}}$$

$$\rightarrow \int e^{-\frac{t}{10}} \cos(t) dt = \frac{1}{101} (-10 \cos(t) e^{-\frac{t}{10}} + 100 \sin(t) e^{-\frac{t}{10}})$$

$$\rightarrow z = e^{\frac{t}{10}} \int e^{-\frac{t}{10}} (\cos(t)) dt = e^{\frac{t}{10}} (\frac{1}{101} (-10\cos(t)) e^{-\frac{t}{10}} + 100\sin(t) e^{-\frac{t}{10}}) + c$$
$$= \frac{10}{101} (10\sin(t) - \cos(t)) + c e^{\frac{t}{10}}$$

$$z = \frac{10}{101} \left(10sin(t) - cos(t) \right) + ce^{\frac{t}{10}} \text{ but } y = z^{-1} \rightarrow z = y^{-1}$$
$$z = \frac{10}{101} \left(10sin(t) - cos(t) \right) + ce^{\frac{t}{10}} \rightarrow y^{-1}$$
$$= \frac{10}{101} \left(10sin(t) - cos(t) \right) + ce^{\frac{t}{10}}$$
$$\rightarrow y = \frac{1}{\frac{10}{101} \left(10sin(t) - cos(t) \right) + ce^{\frac{t}{10}}}$$

To verify the solution:

$$y = \frac{1}{\frac{10}{101}(10sin(t) - cos(t)) + ce^{\frac{t}{10}}} \rightarrow y' = \frac{-\frac{10}{101}(10cos(t) + sin(t)) - \frac{1}{10}ce^{\frac{t}{10}}}{(\frac{10}{101}(10sin(t) - cos(t)) + ce^{\frac{t}{10}})^2}$$
$$\rightarrow y' = \frac{-\frac{10}{101}(10cos(t) + sin(t)) - \frac{1}{10}ce^{\frac{t}{10}}}{\frac{1}{y^2}} =$$
$$\frac{-\frac{1}{10}(\frac{10}{101}(10sin(t) - cos(t)) + ce^{\frac{t}{10}}) + \frac{101}{101}cos(t)}{\frac{1}{y^2}}$$
$$\rightarrow y' = \frac{-\frac{1}{10}(\frac{1}{y}) + cos(t)}{\frac{1}{y^2}} = (-\frac{1}{10}(\frac{1}{y}) + cos(t))y^2$$

→
$$y' = -\frac{1}{10}y + \cos(t)y^2$$
 correct (\textcircled{i})

Exercises

1- Determine if either of the following equations are separable.

$$y' = \cos(t+y) + \cos(t-y), \qquad y' = \cos(t+y) + \sin(t-y),$$

2- Solve the following homogeneous equations

(

(i)
$$y' = \frac{y}{t}(\frac{y}{t} + 1),$$

(ii) $y' = \frac{t^2 - 3y^2}{ty},$
(iii) $y' = \frac{3t - 4y}{3t + 4y},$
(iv) $ty' = y + \tan(\frac{y}{t}),$
(v) $y' = \frac{y}{t}[\frac{1}{\ln(\frac{y}{t})} - 1].$

3- Determine which of the following differential equations are exact

$$(t^{2} + ty)dt + tydy = 0,$$

$$(2y + y^{2})dt - tdy = 0,$$

$$t^{2}y^{3}dt + t^{3}y^{2}dy = 0,$$

$$t^{e^{t}} + y)dt + (2y + t + ye^{y})dy = 0.$$

4- Show that the following equations are exact and then solve them

$$\begin{aligned} \frac{xdx}{(x^2+y^2)^{3/2}} + \frac{ydy}{(x^2+y^2)^{3/2}}, \\ \frac{dy}{dx} &= \frac{y+6x^2}{x(2-\ln(x))}, \\ tdt + ydy &= 0. \end{aligned}$$

, 5- Solve the IVP,

$$y' + y = \cos t,$$
 $y(0) = 1.$
 $ty' + 2y = e^t,$ $t > 0,$
 $y' - \frac{3}{t}y = 4y^{-5},$ $y(1) = 2.$