

Ministry of Higher Education and Scientific Research
University of Baghdad
College of Science
Department of Biology



Pathogenic Bacteria


2022-2023

المرحلة الرابعة - الدراساتين الصباحية والمسائية
الفصل الدراسي الاول

تدريسي المادة :

أ.د. رسمية عبد ابو ريشة
م.د. جنان محمد حسن

أ.د. حارث جبار فهد
أ.د. حلا مؤيد رديف



PATHOGENIC BACTERIA: AN OVERVIEW

Lecture one



BY
DR. HARITH J. FAHAD
Professor of pathogenic bacteria

Lecture one: Pathogenic bacteria: an overview

Introduction

The science of medical microbiology dates back to the pioneering studies of Pasteur and Koch. The methods they developed lead to the first golden age of microbiology (1875-1910), when many bacterial diseases and the organisms responsible for them were defined.

In the first half of the 20th century, scientists studied the structure, physiology, and genetics of microbes in detail and began to answer questions relating to the links between specific microbial properties and disease.

Contributors

Louise Pasteur (1822-1895):

- A French scientist, considered as the **father of microbiology**.
- In 1862, he proposed **germ theory** of disease.
- In 1867, he devised the process of destroying bacteria known as **pasteurization**.
- In 1881, he developed anthrax vaccine and resolved the pebrine silkworm disease.
- In 1885, he developed rabies vaccine.

Robert Koch (1843-1910):

- A German physician, he perfected many bacteriological techniques and known as father of practical bacteriology.
- in 1876, he isolated anthrax bacilli from infected cattle in a pure culture.
- In 1882, he isolated the *Mycobacterium tuberculosis* that causes tuberculosis.
- in 1883 he isolated *Vibrio cholerae* the causative agent of cholera.
- In 1883, he verified the germ theory of disease by relating a specific organism to specific disease.

Lecture one: Pathogenic bacteria: an overview

- In 1884, he established scientific bases called **Koch's postulates**:
 - 1- The organism should be present in all animals suffering from the disease and absent in all healthy animals.
 - 2- The suspected organism must be isolated and grown in a pure culture.
 - 3- The same disease must result when the isolated microorganism is inoculated into a healthy host.
 - 4- The same organism must be isolated again from the experimentally infected animal and shown to be identical with the original isolate.

Joseph Lister (1827-1912):

A British surgeon first introduced the technique to reduce microbes in medical settings and prevent wound infections in 1867. He always known as the father of antiseptic surgery or father of modern surgery.

Host parasite relationship

From the moment of birth, human beings are continuously exposed to microorganisms including bacteria. Some of the interactions between human and microorganisms are essential for the wellbeing of individual but others result in disease.

A parasite is an organism that reside in or within another living organism in order to find the environment, nutrient that required for the growth and reproduction.

The surface of tissue of the skin, oral cavity, respiratory tract, gastrointestinal tract, genital and urinary tract are populated by bacteria and another microorganism.

Microbial flora (microbiota) are microbial population frequently found in association with particular tissue in normal healthy individuals.

Lecture one: Pathogenic bacteria: an overview

Transient flora are acquired from environment and established themselves briefly but tend to be excluded by competition from resident or by the host innate or immune defense mechanisms. They found in association with skin and mucous membranes, they may be present for hours, days, or few weeks. They may or may not be pathogenic.

Resident flora are strains that have an established niche at one of the many body sites; which they occupy indefinitely. They found regularly inside the body (endosymbiosis) or on surface (ectosymbiosis) usually not pathogenic but may be opportunistic.

Normal flora of the skin

- *Staphylococcus epidermidis*
- *Staphylococcus aureus* (in small numbers)
- Nonpathogenic *Neisseria* species
- Alpha-hemolytic and nonhemolytic streptococci
- Diphtheroids
- *Propionibacterium* species
- *Peptostreptococcus* species
- Small numbers of other organisms (*Candida* species, *Acinetobacter* species, etc)

Normal flora of the mouth

- *Staphylococcus* spp.
- *Neisseria* spp.
- *Lactobacillus* spp.
- *Bacteroides* spp.
- The mucous membrane of mouth and pharynx are often sterile at birth, after birth some viridans streptococci become established.

Lecture one: Pathogenic bacteria: an overview

Normal flora of upper respiratory tract

- Alpha-hemolytic and nonhemolytic streptococci
- *Haemophilus*.
- Pneumococci.
- Mycoplasma.
- *Bacteroides* spp.

Normal flora of eye

- Staphylococci.
- Streptococci.
- *Corynebacterium xerosis*.

Normal flora of ears

- Staphylococci.
- Streptococci.
- *Bacillus* spp.

Normal flora of gastrointestinal tract: Upper intestine

- Lactobacilli.
- Enterococci.

Normal flora of gastrointestinal tract: Colon

- 96-99% *anaerobic bacteria*
- Lactobacilli.
- Streptococci.
- *Fusobacterium* spp.
- *Bifidobacterium* spp.
- *Bacteriodes* spp.
- *Clostridium perfringens*.

Lecture one: Pathogenic bacteria: an overview

- Enterobacteriaceae.
- Small numbers of *Proteus* and *Pseudomonas*.

The important functions of intestinal micro biota

The important functions of intestinal microbiota can be divided into three major categories:

- 1- Protective functions: in which the resident bacteria displace and inhibit microbial pathogens indirectly by competing for nutrient and receptors or directly through the production of antimicrobial factors, such as bacteriocins and lactic acid.
- 2- The development and function of the mucosal immune system: such as induce the secretion of IgA.
- 3- Metabolic functions: Such as synthesize vitamin K, biotin, and folate and enhance ion absorption certain bacteria metabolize dietary carcinogens and assist with fermentation of non-digestible dietary residue.

Pathogenicity of microorganisms

Pathogenicity is the capacity of microorganisms to cause disease. Pathogens are microorganisms capable of causing disease. They are of two types:

- 1- **Opportunistic pathogen:** agents capable of causing disease only when the host resistance is impaired (when the patient is compromised) or they are normal flora cause disease when the immune response of the host is suppressed. For example, *Clostridium perfringens* is normal flora in colon; however, it causes gas gangrene in locally damaged tissues.
- 2- **True pathogen:** they possess properties that enable them to overcome the body defenses and infect the tissues of normal healthy subject.

Lecture one: Pathogenic bacteria: an overview

A few substances play role in production of disease.

1- Toxins

Microbial toxins are usually grouped as endotoxin and exotoxin. Exotoxins are proteins that are most often excreted from the cell. However, some exotoxins accumulate inside the cell and are either injected directly into the host or are released by cell lysis. Endotoxins are lipid molecules that are components of the bacterial cell membrane. The essential features of each group are listed below:

Table 1: Essential features of exotoxins and endotoxins

Exotoxins	Exotoxins
Excreted by living cell; high concentrations in liquid medium	Integral part of the cell wall of Gram-negative bacteria; released on bacterial death and in part during growth; may not need to be released to have biologic activity
Produced by both Gram-positive and Gram-negative bacteria	Found only in Gram-negative bacteria
Polypeptides with a molecular weight of 10,000–900,000	Lipopolysaccharide complexes; lipid A portion probably responsible for toxicity
Relatively unstable; toxicity often destroyed rapidly by heating at temperatures above 60°C	Relatively stable; withstand heating at temperatures above 60°C for hours without loss of toxicity
Highly antigenic; stimulate formation of high-titer antitoxin; antitoxin neutralizes toxin	Weakly immunogenic; antibodies are antitoxic and protective; relationship between antibody titers and protection from disease is less clear than with exotoxins
Converted to antigenic, nontoxic toxoids by formalin, acid, heat, and so on; toxoids are used to immunize (eg, tetanus toxoid)	Not converted to toxoids
Highly toxic; fatal to animals in microgram quantities or less	Moderately toxic; fatal for animals in tens to hundreds of micrograms
Usually bind to specific receptors on cells	Specific receptors not found on cells
Usually do not produce fever in the host	Usually produce fever in the host by release of interleukin-1 and other
Frequently controlled by extrachromosomal genes (eg, plasmids)	Synthesis directed by chromosomal genes

2. Extracellular enzymes

Many species of bacteria produce enzymes that are not intrinsically toxic but do play important role in the infectious process.

A- Tissue-degrading enzymes

Many species of bacteria produce tissue-degrading enzymes. The best characterized are enzymes from *C. perfringens*, and to a lesser extent, the aerobic bacteria, *S. aureus* and group A streptococci.

Lecture one: Pathogenic bacteria: an overview

Collagenase: a proteolytic enzyme produced by *C. perfringens*, which degrades collagen, the major protein of fibrous connective tissue, and promotes spread of infection in tissue.

Coagulase produced by *S. aureus*, which works in conjunction with blood factors to coagulate plasma. Coagulase contributes to the formation of fibrin walls around staphylococcal lesions, which helps them persist in tissues.

Hyaluronidases are enzymes that hydrolyze hyaluronic acid, a constituent of the ground substance of connective tissue. They are produced by many bacteria (e.g., staphylococci, streptococci, and anaerobes) and aid in their spread through tissues.

Streptokinase (fibrinolysin) produced by hemolytic streptococci activates a proteolytic enzyme of plasma. This enzyme is then able to dissolve coagulated plasma and probably aids in the rapid spread of streptococci through tissues.

Cytolysins: they dissolve red blood cells (**Hemolysins**) or kill tissue cells or leukocytes (**Leucocidins**). Many bacteria produce these substances such as Clostridia and staphylococci. Most gram negative rods isolated from sites of disease produce hemolysins. For example, *E. coli* strains that cause UTI typically produce hemolysins, strains that are part of the normal gastrointestinal flora may or may not produce hemolysins.


B. IgA1 proteases

Some bacteria that cause disease produce IgA1 proteases. IgA1 protease is an important virulence factor of the pathogens; *N. gonorrhoeae*, *N. meningitidis*, *H. influenzae*, and *S. pneumoniae*.

The enzymes are also produced by some streptococci associated with dental disease, and a few strains of other species that occasionally cause disease production of IgA1 protease allows pathogens to inactivate the primary

Lecture one: Pathogenic bacteria: an overview

antibody found on mucosal surfaces and thereby eliminates protection of the host by the antibody.



PATHOGENESIS OF BACTERIAL INFECTIONS

Lecture two



BY
DR. HARITH J. FAHAD
Professor of pathogenic bacteria

Pathogenesis

Infectious diseases are complex and involve a series of complex and shifting interactions between the invading organism and the host these interactions include the following:

1. The organism's ability to breach host barriers and to evade destruction by innate local and tissue host defenses.
2. The organism's biochemical tactics to replicate, to spread, to establish infection, and to cause disease.
3. The microbe's ability to transmit to a new susceptible host.
4. the body's innate and adaptive immunologic ability to control and eliminate the invading parasite.

Infection may imply colonization, multiplication, invasion or persistence of a pathogen on or within a host.

Infectious disease is used to describe an infection that causes significant overt damage to the host. There are two broad qualities of pathogenic bacteria underlie the means by which they cause disease: invasiveness and toxigenesis.

Invasiveness is the ability to invade tissues. This encompasses mechanisms for colonization (adherence and initial multiplication), ability to bypass or overcome host defense mechanisms, and the production of extracellular substances "invasins" which facilitate the actual invasive process.

Toxigenesis is the ability to produce toxins. Toxic substances may be transported by blood and lymph and cause cytotoxic effects at tissue sites remote from the original point of invasion or growth.

Pathogenesis is the process of causing disease. It depends on the:

1. Immune status of the host
2. Nature of the species or strain (virulence factors)

Lecture two: Pathogenesis of bacterial infections

3. Phase of microbial growth: bacteria in log phase are more likely to overcome host resistance than those in late phases.
4. Number of organisms in the initial exposure (dosage of pathogens); for example *Salmonella enterica* serovar Typhi infection need small dose, while *Salmonella enteritidis* infection need to large does.

Virulence

The term virulence refers to the degree of pathogenicity of the microbe. Virulence is usually expressed as LD₅₀. LD₅₀ is the number of microorganism or quantity of toxin that necessary to kill 50% of the population in a particular time frame.

Virulence might be lost by the followings:

- Attenuation: it can be achieved by successive transfer of pathogen on culture media.
- Loss of capsule: as in *S. pneumoniae*.
- Colony variation; conversion from smooth to rough, for example *Salmonella* and *Shigella*.
- Loss of temperate phage: as in *C. diphtheriae*.
- Heat and desiccation.

Enhancement of virulence by:

- Successive passage in lab animals.
- Presence with other bacteria; for example, *Corynebacterium diphtheriae* and *Streptococcus pyogenes*.

Virulence can be conserved by:

- Lyophilization.

Lecture two: Pathogenesis of bacterial infections

- Using enrichment culture media contain materials such as blood and store in a dark at low temperature.
- Spore forming bacteria resist adverse conditions; hence they have permanent virulence.

Virulence differs according to host factors:

- Age.
- Malnutrition.
- Drugs addiction.
- Metabolic diseases such as diabetes.
- Haemological diseases.
- Immune deficiency.
- Sex.
- Race

Types of diseases

- **Local:** a disease that is restricted to a certain area in the body.
- **Focal:** localized site of disease from which bacteria and their products can spread to other parts.
- **Primary:** the initial disease caused by a pathogen.
- **Secondary:** a primary disease complicated with second pathogen. Ex. Pneumonia following primary influenza especially in elderly people.
- **Mixed:** a disease caused by two or more microorganisms.
- **Inapparent:** a disease that doesn't give rise to any detectable manifestation.

Lecture two: Pathogenesis of bacterial infections

- **Latent:** a disease that persists in the tissues in dormant state and later become manifested usually when the host resistance is lowered.
- **Bacteremia:** a condition in which blood serves as site of presence of bacteria, but the bacteria usually cleared from the vascular system with no harmful effect.
- **Septicemia:** a condition in which blood serves as a site of bacterial multiplication as well as a mean of transfer of infectious agents from one site to another.
- **Pyemia:** the presence of pyogenic bacteria like Staphylococcus in the blood as they spread from one site to another.
- **Toxemia:** is the presence of toxins in the blood stream.

The mechanisms of bacterial pathogenicity

- **To cause disease a pathogen must:**

- 1- Gain access to the host (Entry into host).
- 2- Adhere to host tissues (Adhesion).
- 3- Penetrate or evade host defenses (Invasion).
- 4- Survival in the host.
- 5- Damage the host either directly or via accumulation of microbial wastes.

- **Entry into host**

A. mucous membranes (moist mucosa); most common routes for most pathogens:

- 1- Respiratory tract (most common).
- 2- Gastrointestinal tract.
- 3- Urinary /genital tracts.
- 4- Conjunctiva.

B. Skin (keratinized cutaneous membrane):

- Some pathogens infect hair follicles and sweat glands.
- Few can colonize surface.
- Unless broken, skin is usually an impermeable barrier to microbes.

C. parenteral route

Penetrate skin: punctures, infections, bites, cuts, surgery, etc.

Deposit organisms directly into deeper tissues.

D. Many opportunistic pathogens are carried as part of the normal human flora and this acts as a ready source of infection in the compromised host.

Adhesion

It is an essential preliminary to colonization and then penetration through tissues. At the molecular level, adhesion involves surface interactions between specific receptors on the mammalian cell membrane (usually carbohydrates) and ligands (adhesin, usually proteins) on the bacterial surface. The presence or absence of specific receptors on mammalian cells contributes significantly to tissue specificity of infection.

Invasiveness

Invasiveness is the ability of microorganisms to enter host tissue, multiply and spread. It is mediated by a complex array of molecules, often described as “invasins” which act against the host by breaking down defense of the body.

Types of bacterial invasins

- Spreading factors; bacterial enzymes such as hyaluronidase, collagenase, etc.

Lecture two: Pathogenesis of bacterial infections

- Enzymes that cause cell lysis; such lecithinases, phospholipases, hemolysin, etc.
- Staphylococcal coagulase
- Extracellular digestive enzymes; such as proteases, lipases, glycohydrolaseas, nucleasese, etc.
- Toxins with short-range effects related to invasion; such as anthrax toxin some bacteria, e.g. *Corynebacterium diphtheriae* or *Clostridium tetani*, (toxin producers) are non-invasive. Others such as staphylococci and streptococci are moderately invasive; while anthrax and plague bacilli are highly invasive.

Survival in the host

Most successful pathogens possess structural or biochemical features which allow them to resist the main lines of host internal defense against them. i.e. the phagocytic and immune responses of the host for example;

- The poly-D-glutamate capsule of *Bacillus anthracis* protects the organisms against cell lysis by cationic proteins in sera or in phagocytes.
- The outer membrane of Gram-negative bacteria is a formidable permeability barrier that is not easily penetrated by hydrophobic compound such as bile salts which are harmful to the Bactria.
- Pathogenic mycobacteria have a waxy cell wall that resists attack or digestion by most tissue bactericides.
- Intact lipopolysaccharides (LPS) of Gram-negative pathogens may protect the cell from complement-mediated lysis or the action of lysozyme.
- Almost all principal pathogens that cause pneumonia and meningitis have antiphagocytic polysaccharide capsules.

Damage the host

Tissue damage and the manifestations of disease may also result from interaction between the host's immune mechanisms and the invading organisms or its products. Reactions between high concentrations of antibody, soluble microbial antigens, and complement can deposit immune complexes in tissues and cause acute inflammatory reactions and immune complex disease.

Epidemiology

The term epidemiology is derived from the Greek word **epidemios** which means among people.

It is the study of the distribution and determinants of health-related states or events (including diseases), and the application of this study to the control of diseases and other health problems.

Lec 3: Lab technique in epidemiology

For epidemiological investigations the strains of bacteria can be divide into types, which are taxonomic categories below the species level. Various typing methods can be used; biotyping, antibiograms, serotyping, bacteriocin typing and pahge typing.

- 1. Biotyping:** Biochemical identification of bacterial isolates.
- 2. Serotyping:** The use of antigenic differences among bacterial strains. For example, flagellar, cell wall, and capsular antigens are frequently used for *Salmonella*, *Shigella*, and eteropathogenic *E. coli* typing. M protein serotyping is useful in epidemiology of glomerulonephritis caused by *S. pyogenes*. Capsular swelling (Quellung reaction) is used in serotyping of *Klebsiella* spp.
- 3. Antibiogram:** A test employed for the determination of susceptibility of microorganisms to antibiotics or other antimicrobial agents.
- 4. Phage typing:** Using the bacteriophages in identification of bacterial strains. Ex. *S. aureus* has 22 phages for identification.
- 5. Bacteriocin typing:** Bacteriocins are antimicrobial substances produced by bacterial strains. It is used in identification of *Serratia marcescens*, *E. coli*, *P. mirabilis*, and *Shigella* spp.
- 6. Genetic methods:** Direct methods including hybridization of DNA or indirect methods using restricted endonucleases analysis.

Pattern of diseases in community

- **Endemic:** a disease found constantly in low percentage with a certain area.
Ex. Cholera in south east Asia.

- **Epidemic:** the outbreak that happens suddenly in high percentage of disease in susceptible persons within certain population. Ex. meningitis in Iraq in 1976.
- **Pandemic:** communicable infections that are wide spread in a region, sometimes worldwide, and have high attack rates. Ex. Asian influenza.
- **Sporadic** is a disease occurs in an irregular pattern.

Patterns of transmission of diseases

- 1- Horizontal: disease is spread thorough a population from one infected person to another (kissing, sneezing)
- 2- Vertical: the disease is transmitted from parent to offspring (ovum, sperm, placenta, milk)
- 3- Contact transmission:
 - Direct – kissing, sex
 - Droplets – talking
 - Vertical – mother to fetus
 - Vector
- 4- Indirect contact:
 - Food, water, and biological products (blood, serum , tissue)
 - Fomite (door knobs, toilet seats, etc.)
 - Air
 - Droplet nuclei (dried microscopic residue)
 - Aerosols (dust or moisture particles)

Frequency measures

- **Prevalence** is the total number of cases in a population at any given time.

- **Incidence** of disease is the number of new cases over a defined period of time.
- **Mortality rate** is the number of individuals that die as a result of the outbreak of disease within certain period of time.
- **Nosocomial infections:** hospital acquired infections.

Clinical stages of a disease in the host

A- Acute disease

It is characterized by symptoms that usually appears quickly and become very intense; thereafter, subsides when the host immune system overwhelmed the pathogen or its toxic products. Ex. Childhood diseases like measles, mumps, various types of influenza and chicken pox. Acute disease involves three periods:

- 1- Incubation period:** it is the time between exposure to the organism and appearance of the first symptoms of the disease.
- 2- Acute period:** symptoms of the disease are at their peak. ex. Fever and cough in respiratory infections, diarrhea and vomiting in intestinal tract infections.
- 3- Convalescence period:** a period characterized by a sharp decline in symptoms.

B- Chronic diseases and persistence

Their symptoms are expressed over a long period of time. Infectious agent here is an intracellular parasite like *Brucella* and *M. tuberculosis*. Some microorganisms that persist in host give rise to mild symptoms or no symptoms and they are actively shed from the host. The individual shedding these microorganisms called carrier. The microorganism may persist in the host for

days, months, or even years like the causative agents of cholera, typhoid, and diphtheria.

Reservoirs and sources of diseases

It includes animate and inanimate

1. Animate

It includes patients, carriers, and animals.

a. Patients, they represent the most important reservoirs for many viral and bacterial disease like measles, mumps, poliomyelitis, whooping cough and sexually transmitted diseases.

b. Carriers, infected healthy individual has no symptoms (asymptomatic), or very mild form of a disease, yet they both can spread disease to others many bacterial pathogens. Three types of carriers are recognized:

- **A convalescent** carrier is a person who has recovered from the symptoms of an infectious disease but is still capable of transmitting pathogens to others.
- **A healthy** carrier is a person who or animal that harbors a specific infectious agent in the absence of discernible clinical disease and serves as a potential source of infection.
- **A chronic** carrier carries infectious agents for long periods of time.

c. Animals

Zoonoses is any infection or disease that is transmitted to man from animals (a disease occurring primary in animal and transmitted secondary to human) like *Salmonella* is transmitted from cattle and poultry through contaminated food causing salmonellosis. Two classes of arthropods serve as reservoirs and sources of infections:

- Class **Insecta** includes flies, mosquitoes, fleas, and lice.

- Class **Arachnida** involves ticks and mites.


The relationship between infectious diseases and arthropods are:

- **Mechanical**, arthropods carry infectious agents on their appendages and not involved in their life cycle. Ex. Flies and *Salmonella*.
- **Biological**: arthropods serve as hosts and reservoirs of the microbial agents. Ex. mosquitoes and malaria parasites.

2: Inanimate

Inanimate sources and reservoirs include:

- **Soil** (*Clostridium tetani* and *Clostridium perfringens*).
- **Water** (*Vibrio cholerae*).
- **Food** (*Salmonella enterica* serovar Typhi, *Clostridium botulinum* and *Campylobacter*).



ENTEROBACTERIACEAE

Lecture 4



PROF. DR. HARITH JABBAR FAHAD
Dept. of Biology, College of Science, University of Baghdad

Enterobacteriaceae

They are large heterogeneous group of aerobes and facultative anaerobes, Gram-negative rods, ferment a wide range of carbohydrates, possess a complex antigenic structure, and produce a variety of bacteriocins such as colicins, toxins (endotoxins and exotoxins) and other virulence factors. The natural habitat is the intestinal tract of humans and animals.

Bacteriocins: proteins are produced by certain strains of bacteria active against some other strains of the same or closely related species.

This family includes 42 genera (*Escherichia*, *Shigella*, *Salmonella*, *Enterobacter*, *Klebsiella*, *Serratia*, *Proteus*, and others). The clinically significant enterobacteriaceae comprises 20-25 species, and other species are encountered infrequently.

Antigenic structure

More than 150 different heat-stable somatic O (lipopolysaccharide) antigens.

More than 100 heat-labile K (capsular) antigens.

More than 50 H (flagellar) antigens.

Diseases caused by enterobacteriaceae

A. *Escherichia coli*

1. Urinary tract infections

Escherichia coli is the most common cause of urinary tract infection and accounts for approximately 90% of first urinary tract infections in young women.

2. *E. coli*-associated diarrheal disease

- Enteropathogenic *E. coli* (EPEC) is an important cause of diarrhea in infants, especially in developing countries. The duration of the EPEC diarrhea can be shortened and the chronic diarrhea cured by

antibiotic treatment. Care in the selection and consumption of foods potentially contaminated with ETEC is highly recommended to help prevent traveler's diarrhea. Antibiotic treatment effectively shortens the duration of disease.

- Enterotoxigenic *E. coli* (ETEC) is a common cause of "traveler's diarrhea" and diarrhea in infants in developing countries. Some strains produce a heat-labile exotoxin (LT) and some produce the heat-stable enterotoxin (STa).
- Shiga toxin-producing *E. coli* (STEC) they produce two cytotoxic toxins, Shiga-like toxin 1 and Shiga-like toxin 2 of the *E. coli* serotypes that produce Shiga toxin, O157:H7 is the most common. STEC has been associated with hemorrhagic colitis, a severe form of diarrhea which can be prevented by thoroughly cooking ground beef and avoiding unpasteurized products such as apple cider.
- Enteroinvasive *E. coli* (EIEC) produces a disease very similar to shigellosis. The disease occurs most commonly in children in developing countries and in travelers to these countries.
- Enteroaggregative *E. coli* (EAEC) causes acute and chronic diarrhea in persons in developing countries, food-borne illnesses in industrialized countries, and have been associated with traveler's diarrhea and persistent diarrhea in patients with HIV.

3. Sepsis

When normal host defenses are inadequate, *E. coli* may reach the bloodstream and cause sepsis.

4. Meningitis

Approximately 80% of *E. coli* from meningitis cases have the K1 antigen. Diagnostic Laboratory Tests.

B. Klebsiella

Klebsiella pneumoniae is present in the nasopharynx and feces of about 5% of normal individuals.

C. Serratia

Serratia marcescens is a common opportunistic pathogen in hospitalized patients. *Serratia* (usually nonpigmented) causes pneumonia, bacteremia, and endocarditis.

D. Proteus

Proteus species produce infections only when the bacteria leave the intestinal tract. They are found in urinary tract infections and produce bacteremia, pneumonia, and focal lesions in debilitated patients or those receiving contaminated intravenous infusions. *P. mirabilis* causes urinary tract infections and occasionally other infections. *P. vulgaris* is important nosocomial pathogen.

Immunity: specific antibodies develop in systemic infections, but it is uncertain whether significant immunity to the organism follows.

Treatment

No single specific therapy is available. The sulfonamides, ampicillin, cephalosporins, fluoroquinolones, and aminoglycosides have marked antibacterial effects against the enterics. However, variation in susceptibility is great, and laboratory tests for antibiotic susceptibility are essential. Multiple drug resistance is common and is under the control of transmissible plasmids.

Epidemiology, Prevention, and Control

Some of the enterics constitute a major problem in hospital infection; these bacteria commonly are transmitted by personnel, instruments, or parenteral medications. Their control depends on handwashing, rigorous

Lecture 4: Enterobacteriaceae

asepsis, sterilization of equipment, disinfection, restraint in intravenous therapy, and strict precautions in keeping the urinary tract sterile (ie, closed drainage).

E. The shigellae

Antigenic Structure

There are more than 40 serotypes.

Pathogenesis and Pathology

Shigella infections (shigellosis) are almost always limited to the gastrointestinal tract; bloodstream invasion is quite rare. Shigellae are highly communicable; the infective dose is low, on the order of 10^2 organisms. The pathogenic species of *Shigella* are *S. dysenteriae*, *S. flexneri*, *S. boydii* and *S. sonnei*.

Toxins

A. Endotoxin

B. *Shigella dysenteriae* Exotoxin

Shigella dysenteriae type 1 (Shiga bacillus) produces a heat-labile exotoxin that affects both the gut (acting as an enterotoxin) and the central nervous system (acting as neurotoxin). The exotoxin is a protein that is antigenic and lethal for experimental animals.

Clinical Findings

After a short incubation period (1–4 days), there is a sudden onset of abdominal pain, fever, and watery diarrhea. In more than half of adult cases, fever and diarrhea subside spontaneously within 2–5 days. However, in children and elderly adults, loss of water and electrolytes may lead to dehydration, acidosis, and even death. The illness caused by *S. dysenteriae* may be particularly severe. On recovery, most persons shed dysentery bacilli

Lecture 4: Enterobacteriaceae

for only a short period, but a few remain chronic intestinal carriers and may have recurrent bouts of the disease. Upon recovery from the infection, most persons develop circulating antibodies to shigellae, but these do not protect against reinfection.

Immunity

Injection of killed shigellae stimulates production of antibodies in serum but fails to protect humans against infections.

Treatment

Ciprofloxacin, ampicillin, doxycycline, and trimethoprim-sulfamethaxazole.

Epidemiology, Prevention, and Control

Shigellae are transmitted by “food, fingers, feces, and flies” from person to person. Most cases of *Shigella* infection occur in children younger than 10 years of age. Because humans are the main recognized host of pathogenic shigellae, control efforts must be directed at eliminating the organisms from this reservoir by:

- (1) sanitary control of water, food, and milk; sewage disposal and fly control;
- (2) isolation of patients and disinfection of excreta;
- (3) detection of subclinical cases and carriers, particularly food handlers;
and
- (4) antibiotic treatment of infected individuals.

F. The *Salmonella*-Arizona group

Salmonellae are often pathogenic for humans and animals when acquired by the oral route. They are transmitted from animals and animal products to humans where they cause enteritis, systemic infection, and enteric fever.

Classification

Currently, the genus *Salmonella* is divided into two species; *Salmonella enterica* (contains subspecies; I, II, IIIa, IV, and VI) and *Salmonella bongori*. Most human illness is caused by *S. enterica* subspecies I strains, written as *S. enterica* subspecies *enterica*. There are more than 2500 serotypes (serovars) of *Salmonella*, including more than 1400 in DNA hybridization group I that can infect human. Four serotypes of salmonellae that cause enteric fever can be identified. They are as follows: *Salmonella* Paratyphi A (serogroup A), *Salmonella* Paratyphi B (serogroup B), *Salmonella* Choleraesuis (serogroup C1), and *Salmonella* Typhi (serogroup D).

Pathogenesis

Salmonella Typhi, *Salmonella* Choleraesuis, and perhaps *Salmonella* Paratyphi A and *Salmonella* Paratyphi B are primarily infecting humans. The organisms almost always enter via the oral route, usually with contaminated food or drink. The mean infective dose to produce clinical or subclinical infection in humans is 10^5 – 10^8 salmonellae (but perhaps as few as 10^3 *S. Typhi* organisms). Among the host factors that contribute to “resistance” to *Salmonella* infection are gastric acidity, normal intestinal microbiota, and local intestinal immunity. Salmonellae produce three main types of disease in humans, but mixed forms are frequent.

A) The "Enteric Fevers" (Typhoid Fever)

This syndrome is produced by only a few of the salmonellae, of which *S. Typhi* (typhoid fever) is the most important. The organisms multiply in intestinal lymphoid tissue and are excreted in stools. After an incubation period of 10–14 days, fever, malaise, headache, constipation, bradycardia, and myalgia occur. The spleen and liver become enlarged.

B) Bacteremia with Focal Lesions

Lecture 4: Enterobacteriaceae

This is associated commonly with *S. Choleraesuis* but may be caused by any *Salmonella* serotype.

C) Enterocolitis

This is the most common manifestation of *Salmonella* infection. Enterocolitis can be caused by any of the more than 1400 group I serotypes of salmonellae. Eight to 48 hours after ingestion of salmonellae, there is nausea, headache, vomiting, and profuse diarrhea, with few leukocytes in the stools. Low-grade fever is common, but the episode usually resolves in 2–3 days.

Treatment

Ampicillin, trimethoprim-sulfamethoxazole, or a third-generation cephalosporin.

Epidemiology

The feces of people who have unsuspected subclinical disease or who are carriers are a more important source of contamination than frank clinical cases that are promptly isolated, such as when carriers working as food handlers are “shedding” organisms. Many animals, including cattle, rodents, and fowl, are naturally infected with a variety of salmonellae and have the bacteria in their tissues (meat), excreta, or eggs. The high incidence of salmonellae in commercially prepared chickens has been widely publicized.

A. Carriers


After manifest or subclinical infection, some individuals continue to harbor salmonellae in their tissues for variable lengths of time (i.e., convalescent carriers or healthy permanent carriers). Three percent of survivors of typhoid fever become permanent carriers, harboring the organisms in the gallbladder, biliary tract, or, rarely, the intestine or urinary tract.

B. Sources of Infection

1. Water, milk and other dairy products (ice cream, cheese, custard).
2. Shellfish, meats and meat products.
3. Dried or frozen eggs.
4. “Recreational” drugs.
5. Animal dyes
6. Household pets.

C. Prevention and Control

1. Sanitary measures must be taken to prevent contamination of food and water by rodents or other animals that excrete salmonellae.
2. Infected poultry, meats, and eggs must be thoroughly cooked.
3. Carriers must not be allowed to work as food handlers or in food preparation areas, and should observe strict personal hygienic precautions.
4. Vaccination is recommended for travelers to endemic regions.



VIBRIO, CAMPYLOBACTER, AND HELICBACTER

Lecture 4 and 5



BY

DR. HARITH J. FAHAD

Professor of pathogenic bacteriology

Lecture 4: Vibrio, Campylobacter, and Helicobacter

Vibrio, Campylobacter, and Helicobacter

Vibrio, *Campylobacter*, and *Helicobacter* are Gram-negative rods that are all widely distributed in nature. The vibrios are found in marine and surface waters. The campylobacters are found in many species of animals, including many domesticated animals. *Vibrio cholerae* produces an enterotoxin that causes cholera, a profuse watery diarrhea that can rapidly lead to dehydration and death. *Campylobacter jejuni* is a common cause of enteritis in humans. *Helicobacter pylori* is associated with gastritis and duodenal ulcer disease.

The Vibrios

Vibrios are among the most common bacteria in surface waters worldwide. They are curved aerobic rods and are motile, possessing a polar flagellum. *V. cholerae* serogroups O1 and O139 cause cholera in humans, and other vibrios, may cause sepsis, or enteritis. The medically important vibrios are listed in Table 1.

Table 1: The Medically Important Vibrios


Organism	Human Disease
<i>V. cholerae</i> serogroups O1 and O139	Epidemic and pandemic cholera
<i>V. cholerae</i> serogroups non-O1/non-O139	Cholera-like diarrhea; mild diarrhea; rarely, extraintestinal infection
<i>V. parahaemolyticus</i>	Gastroenteritis, wound infections, septicemia
<i>V. vulnificus</i>	Gastroenteritis, wound infections, septicemia

Vibrio cholerae

V. cholera has at least 206 O antigen groups. The *V. cholerae* serogroup O1 antigen has determinants that make possible further subtyping; these serotypes are Ogawa, Inaba, and Hikojima. Furthermore, two biotypes of

Lecture 4: Vibrio, Campylobacter, and Helicobacter

epidemic *V. cholerae* have been defined, classic and El Tor. The El Tor biotype tends to cause milder disease than the classic biotype.



***Vibrio cholerae* Enterotoxin**

V. cholerae produce a heat-labile enterotoxin with a molecular weight (MW) of about 84,000, consisting of subunits A and B. Ganglioside GM₁ serves as the mucosal receptor for subunit B, which promotes entry of subunit A into the cell. Activation of subunit A₁ yields increased levels of intracellular cyclic adenosine monophosphate (cAMP) and results in prolonged hypersecretion of water and electrolytes. There is increased sodium-dependent chloride secretion, and absorption of sodium and chloride by the microvilli is inhibited (Figure 1). Electrolyte-rich diarrhea occurs with as much as 20–30 L/day, resulting in dehydration, shock, acidosis, and death.

Pathogenesis & Pathology

A person with normal gastric acidity may have to ingest:

- $\geq 10^{10}$ *V. cholera* when the vehicle is water
- $10^2 - 10^4$ organisms when the vehicle is food.

Lecture 4: Vibrio, Campylobacter, and Helicobacter

Cholera is not an invasive infection. Virulent *V. cholerae* organisms attach to the microvilli of the brush border of epithelial cells. There they multiply and liberate cholera toxin and perhaps mucinases and endotoxin.




Figure 1: Mode of action of cholera enterotoxin

Clinical findings

About 50% of infections with classic *V. cholerae* are asymptomatic, as are about 75% of infections with the El Tor biotype. The incubation period after ingestion of a sufficiently high infectious dose of *V. cholerae* is 12 hours to 3 days for persons who develop symptoms, depending largely on the size of the inoculum ingested. Stools, which resemble "rice water," contain mucus, epithelial cells, and large numbers of vibrios. There is rapid loss of fluid and electrolytes, which leads to profound dehydration, circulatory collapse, and anuria. The mortality rate without treatment is between 25% and 50%.

Immunity

Lecture 4: Vibrio, Campylobacter, and Helicobacter

Gastric acid provides some protection against vibrios, including *V. cholerae*. An attack of cholera is followed by immunity to reinfection, but the duration and degree of immunity are not known.

Treatment

- The most important part of therapy consists of water and electrolyte replacement to correct the severe dehydration and salt depletion.
- Many antimicrobial agents are effective against *V. cholerae*. Oral tetracycline and doxycycline.
- Erythromycin and/or azithromycin are an appropriate choice of antimicrobial therapy in children and in pregnant women.

Epidemiology

- 1817 – 1993; eight pandemics of cholera were occurred by *V. cholera* O1 of the classic and El Tor biotype and serotype O139 in Asia, Middle East, Africa, south America and Central America.
- The disease is rare in North America.
- Cholera is endemic in India and in Southeast Asia.
- Cholera is a disease that is spread by:
 1. Contact involving individuals with mild or early illness.
 2. Water, food, and flies.
- Only 1–5% of exposed susceptible persons develop disease.

Prevention and Control

- Infection control rests on education and on improvement of sanitation, particularly, food and water.
- Patients should be isolated, their excreta disinfected, and contacts followed up.
- Chemoprophylaxis with antimicrobial drugs may have a place.

Lecture 4: Vibrio, Campylobacter, and Helicobacter

- Repeated injection of a vaccine containing either lipopolysaccharides extracted from vibrios or dense *Vibrio* suspensions can confer limited protection to heavily exposed persons (e.g. family contacts) but is not effective as an epidemic control measure.

Campylobacters

Campylobacters cause both diarrheal and systemic diseases, and are among the most widespread causes of infection, worldwide. *C. jejuni* and the other campylobacters are Gram-negative rods with comma, S, or “gull wing” shapes (Figure 2). They are motile, with a single polar flagellum, and do not form spores. *C. jejuni* is a very common cause of diarrhea in humans.

The campylobacters have:

- LPS with endotoxic activity.
- Cytopathic extracellular toxins.
- Enterotoxins.

The significance of the toxins in human disease is not well defined.



Figure 2: Shapes of *Campylobacter*

Pathogenesis and Pathology

The source of infection may be:

Lecture 4: Vibrio, Campylobacter, and Helicobacter

1. Food and drink (e.g. milk, undercooked fowl).
2. Contact with infected animals (especially poultry) or humans and their excreta.

C. jejuni is susceptible to gastric acid, and ingestion of about 10^4 organisms is usually necessary to produce infection. The organisms multiply in the small intestine, invade the epithelium, and produce inflammation that results in the appearance of red and white blood cells in the stools. Occasionally, the bloodstream is invaded, and a clinical picture of enteric fever develops.

Clinical findings

Clinical manifestations of *C. jejuni* gastroenteritis are acute onset of crampy abdominal pain, profuse diarrhea that may be grossly bloody, headache, malaise, and fever. Usually, the illness is self-limited to a period of 5–8 days, but occasionally it continues for longer times.

Helicobacter pylori

H. pylori is a spiral-shaped (Figure 3), Gram-negative rods. *H. pylori* is associated with antral gastritis, duodenal (peptic) ulcer disease, gastric ulcers, gastric adenocarcinoma, and gastric mucosa-associated lymphoid tissue (MALT) lymphomas. *H. pylori* is a major risk for gastric cancer.



Figure 3: Helicobacter pylori

Pathogenesis and pathology

H. pylori grows optimally at a pH of 6.0–7.0, it would be killed or not grow at the pH within the gastric lumen (pH 1–3). Gastric mucus is relatively impermeable to acid and has a strong buffering capacity. On the lumen side of the mucus, the pH is low (1.0–3.0); on the epithelial side, the pH is about 5.0–7.0. *H. pylori* is found deep in the mucous layer near the epithelial surface where a near physiologic pH is present. *H. pylori* also produces a protease that modifies the gastric mucus and further reduces the ability of acid to diffuse through the mucus. *H. pylori* produces potent urease activity, which yields production of ammonia and further buffering of acid. *H. pylori* is quite motile, even in mucus, and is able to find its way to the epithelial surface (Figure 4).




Figure 4: *Helicobacter pylori* pathogenesis

Clinical findings

Acute infection can yield an upper gastrointestinal illness with nausea and pain; vomiting and fever may also be present. The acute symptoms may last for less than 1 week or as long as 2 weeks. After colonization, *H. pylori* infection persists for years, perhaps decades, or even a lifetime. About 90% of patients with duodenal ulcers and 50–80% of those with gastric ulcer have *H. pylori* infection.


Treatment

Triple therapy with metronidazole and either bismuth subsalicylate or bismuth subcitrate plus either amoxicillin or tetracycline for 14 days eradicates *H. pylori* infection in 70-95% of patients. An acid-suppressing agent given for 4 to 6 weeks enhances ulcer healing.

Epidemiology

H. pylori is present on the gastric mucosa of:

- Fewer than 20% of persons younger than 30 years old.
- 40–60% of people age 60 years.
- In developing countries, most children are infected by the age of 10, and prevalence remains high (up to 90%) among adults of all age groups.



LECTURE 7: SPORE FORMER GRAM POSITIVE BACILLI

Lecture seven



BY
Dr. Harith Jabbar Fahad

I- *Bacillus*

Saprophytes found in soil, water, air and vegetation. The mostly important genera are *B. anthracis*, *B. cereus*, *B. thuringiensis*, *B. subtilis* and *B. megaterium*.

B. anthracis

General characteristics

Members of this species are non-motile, facultative anaerobes, large bacilli measuring 4-8 μm in length and 0.5-1.0 μm in width. They have square ends and arranged in long chains, usually encapsulated with centric spore. Colonies on agar plate are large, opaque, white, medusa head, and have “cut glass” appearance in transmitted light.

Antigenic structure

1. Capsular polypeptide (Polyglutamic acid)
2. Somatic polysaccharides
3. Protein exotoxin complex

Pathogenicity

B. anthracis is a pathogen of animals (herbivorous). Spores found in soil and on vegetation ingested, inhaled or gain entrance through abraded skin or mucosa. The spores germinate in the tissue and transformed into vegetative cells, which produce the exotoxin. The capsule inhibits phagocytosis. In human, the disease considered as an occupational hazard.

Diseases in human

This bacteria causes four types of disease; Skin infection (cutaneous anthrax), Wool sorters disease, Septicemia, and Gastrointestinal anthrax.

II- *Clostridium*

Saprophytes found in soil, water, air and vegetation. Gram positive spore forming bacilli. Measuring 3-8 μm in length and 0.4-1.2 μm in width. Obligate anaerobes. Some of them are non-motile. Most of them are unencapsulated. Few species parasitize intestinal tract of human and animals. Produce highly toxic exotoxins.

C. botulinum

General characteristics

Found in soil and occasionally in animal feces. Spore are highly resistant to heat, withstanding 100°C for 3-5 hours. Diminish at acid pH or high salt concentration. During growth and autolysis, the bacterium liberates toxins. Seven distinct antigenic varieties (A-G). Type A toxin is a complex consisting of neurotoxin and hemagglutinin which protect the neurotoxin from stomach acid and enzymes. Probably 1-2 μg is lethal to human.

Pathogenicity

General characteristics

C. botulinum causes botulism “an intoxication resulting from the ingestion of food in which *C. botulinum* has grown and produced toxin”. Spiced, smoked vacuum packed or canned alkaline foods are eaten without cooking. The toxin acts by blocking release of acetylcholine at synapses and neuromuscular joints producing respiratory paralysis. Symptoms begin after 18-36 hrs. after ingestion of toxic food; visual disturbance, inability to swallow, speech difficulty and death occurs from respiratory paralysis or cardiac arrest.

Botulism is a disease of high mortality rate. There is no fever, patient remains fully conscious until short time before death. Those who recover does not develop antitoxin in the blood.

C. perfringens

General characteristics

It is implicated in gas gangrene. *C. perfringens* is indigenous members of the intestinal tract of humans and animals. Saprophytes found in soil. Large encapsulated, non-motile, spore former bacilli, produces eleven toxins. Alpha toxin is the most important (acts as lecithinase). Also, they produce collagenase, hemolysin, proteinase, DNase, and some of them produce enterotoxin.

Disease produced

1. Gas gangrene.
2. Food poisoning
2. Enteritis necroticans
3. Uterine infections

C. tetani

General characteristics


Found in soil, human intestine (25%) and animal feces. Motile bacilli forming tennis racket shaped terminal spores. Obligate anaerobe but can stand certain concentration of oxygen. Can be distinguished by specific flagellar antigen. All share common somatic antigen. It produced two toxins; Tetanospasmine and Tetanolysin

Pathogenesis

Tetanus results from small wounds which contaminated by *C. tetani* spore that germinate and produce toxin. The infection remains localized with minimal inflammatory damage. The toxin produced upon growth, sporulation and lysis of cells. The toxin migrates along neural paths from a local wound to sites of action in the CNS. Severe painful spasms and rigidity of voluntary muscles. The characteristic symptom of lock jaw involves spasms of masseter muscle, followed by progressive rigidity and violent spasms of trunk and limb muscles. Spasms of the pharyngeal muscles cause difficulty in swallowing. Death usually results from interference with mechanisms of respiration. Symptoms appear after 4-6 days to 6 weeks incubation time.

C. difficile

Causes antibiotic associated diarrhea, colitis and pseudomembranous colitis in human. These infections result from overgrowth of *C. difficile* in the colon after the normal flora has been disturbed by antibiotic therapy.



Staphylococci

Lecture 8

Dr. Harith J. Fahad
Professor of pathogenic bacteriology

Gram positive cocci

Genus: *Staphylococcus*

- **General characteristics**

Staphylococci are non sporulating, non-motile Gram-positive cocci. Microscopically, these organisms are typically found in grape-like clusters and tetrads, as well as in pairs and sometimes in short chains. Usually unencapsulated or have limited capsule formation. When grown on blood agar, staphylococci form small (1 to 2 mm), smooth, round colonies that are often pigmented and may be surrounded by a zone of β -hemolysis.

Staphylococci are very hardy organisms and can resist drying, withstand 10% NaCl broth, and will survive at temperatures between 10° and 45° C. Because staphylococci are facultative anaerobes, they will grow in the presence or absence of oxygen. Staphylococci are catalase positive. They reside on mucous membranes, skin and anterior nares.

Classification of species depends mainly on:

1. Aerobic acid production from different carbohydrates.
2. Coagulase activity.
3. Haemolysis.
4. Nitrate reduction.
5. Genetic methods: ribotyping, DNA - DNA hybridization.

Coagulase production separates the staphylococci into **two** major groups:

A- Coagulase positive: The coagulase-positive species *S. aureus*, *S. intermedius*, *S. delphini*, *S. schleiferi* subsp. *coagulans* and the coagulase-variable species *S. hyicus* are regarded as potentially serious pathogens.

B- Coagulase negative:

S. epidermidis, *S. saprophyticus* and many other species.

In general, the staphylococci are variably sensitive to antimicrobial drugs. Resistance could be produced by:

- 1. Production of β - lactamase (penicillinase):** It is controlled by a plasmid, thus they are resistant to penicillins and cephalosporins (more than **95 %** of staphylococci isolates are resistant to those antibiotics).
- 2. Methicillin Resistant *Staphylococcus aureus* (MRSA):** Methicillin is β - lactamase resistant. The responsible genes may be residing in the plasmid or mostly in the chromosome. This character is important especially in nosocomial infection. Bacteria that are resistant to methicillin are also **resistant to all** other antibiotics in the beta-lactam class, which includes penicillin derivatives and cephalosporins. The only antibiotics available to kill MRSA are powerful and potentially toxic options such as **vancomycin**
- 3. Plasmids - mediated resistance** to tetracycline, erythromycin, and aminoglycosides.

I- *Staphylococcus aureus*

❖ Antigenic structure

- 1- Tcichoic acids** (PG linked); **lipoteichoic acids** (membrane associated).
Regulates cationic environment.
- 2- Protein A:** It is responsible for agglutination test known as **COAGGLUTINATION** it combined with Fc portion of IgG molecule. The Fab portion of IgG bound to protein **A** is free to combine with a specific antigen. Thus, it reduces the false positive reactions.
- 3- Capsular polysaccharides.** Eleven serotypes have been reported. Types 1 and 2 are highly encapsulated, mucoid strains that are virulent for experimental animals, but rarely encountered among clinical isolates of

S. aureus. These strains produce a "**microcapsule**" which may be antiphagocytic.

- 4- **Peptidoglycan:** can lead to fever and alternative complement activation that leads to inflammation followed by leukopenia, thrombocytopenia, and shock.
- 5- **Adhesins:** Specific cell wall associated surface proteins of *S. aureus* that bind to matrix proteins such as fibronectin, fibrinogen (clumping factor), collagen, bone sialoprotein, etc. These binding activities are thought to be involved in binding and colonization of various body sites and specific cell types.

❖ **Toxins and enzymes**

1- **Staphylolysins:** exotoxins and they are α , β , γ , δ haemolysins:

- *Alpha- haemolysin;* It is a dermonecrotic protein, dissolves rabbit erythrocytes, damages platelets and has powerful action on vascular smooth muscles.
- *Beta- haemolysin:* Dissolves sheep erythrocytes upon incubation for 1 hr at 37°C, and 18 hr at 10°C (this is called hot and cold reaction). Antigenicity is distinct from other haemolysins. Toxic for many kinds of cells including human erythrocytes.
- *Gamma- haemolysin:* is lytic for erythrocytes from different mammalian species and also cytotoxic for leukocytes. Gamma-toxin has also been proposed to play a role in the pathogenesis of toxic shock syndrome (TSS) together with toxic shock syndrome toxin 1 (TSST-1).
- *Delta- haemolysin:* Dissolves erythrocytes of sheep, mice, guinea pigs, rabbits, horses, rats, as well as human. Leukotoxic as well as dermonecrotic.

2- Leukocidin (Panton-Valentine Leukocidin):

Found in 5% of all *S. aureus* strains and in 50% of ones from abscesses. Lethal to PMNs, disrupts their membranes through pore formation which leads to increased permeability.

3- Enterotoxins:

It is a neurotoxin affects the vomiting center in the CNS, which causes vomiting and diarrhea. The infection needs the presence of the bacteria, while the toxication does not.

4- Toxic shock syndrome toxin-1 (TSST-1):

Found in almost all *S. aureus* strains, isolated from individuals suffering from TSST. About 90% of healthy individual have antibodies against TSST-1. TSST patients either do not have antibodies or have them at very low levels.

5- Exfoliatine toxin (ET):

It is involved in Staphylococcal scalded skin syndrome (SSSS). Mostly phage group II strains of *S. aureus* controlled by a plasmid or chromosomal gene or both.

6- Coagulase:

Plasma clotting protein deposits fibrin on the surface of staphylococci, perhaps altering their ingestion by phagocytic cells or their destruction within such cells. Coagulase is considered as an **invasive** factor rather than spreading factor.

7- Clumping Factor (fibrinogen binding protein):

Clumping factor allows *S. aureus* to adhere to fibrinogen. Coagulase and clumping factor are **distinct** entities.

8- Other enzymes: Such as hyaluronidase, staphylokinase, proteases, lipases, β -lactamase and thermostable nucleases (DNases, RNases).

❖ Pathogenicity**a. Infections of the skin and associated structures:**

Pyoderma (purulent skin infections): abscesses, furuncles, carbuncles, and impetigo.

b. Staphylococcal Scalded Skin Syndrome (SSSS):

Most often observed in infants and young children. The syndrome begins as erythema around the mouth and nose and spreads rapidly to affect the skin of the neck, the trunk, and sometimes the extremities. The epidermal necrolysis mediated by the toxin exfoliation results in extensive areas of denuded skin. The intraepithelial split in **SSSS** is in the granulosal layer.

c. Gastrointestinal tract Staphylococcal diseases:

- i. **Staphylococcal enteritis:** patients receiving chemotherapeutic agents especially via the oral route (reduce the normal flora).
- ii. **Staphylococcal food poisoning.**

d. Deeper infections:

They are either primary infections or may stem from an infection that has metastasized from a cutaneous infection or from carrier site. They are not likely to occur in healthy individuals but rather in those who have a precondition such as extensive surgery, burns, diabetes, cystic fibrosis, lower respiratory tract viral infections, ulcers and immunodeficient (or immunosuppressed) individuals, *S. aureus* can cause infection in many tissues and systems and organs: e.g. endocarditis, cystitis, meningitis, pneumonia, septicaemia, infection of post-operative wounds and others like osteomyelitis

e. **Toxic shock syndrome:** a multisystem, febrile illness with hypotension, vomiting, diarrhea, rash with subsequent palmer and plantar desquamation. Hyperaemia of mucous membranes can occur regularly.

❖ **Diagnostic lab tests**

- a) **Specimens:** it depends on the localization of infection: surface swab, pus, blood, tracheal aspirate, spinal fluid. Antibody determination in serum is rarely of value in the beginning of infection but when there is chronic infection antibodies are very important to be detected.
- b) **Stained smears:** "after taking the specimens" typical staphylococci appear; gram positive bacteria in grape shaped could be seen in stained smear of pus and sputum.
- c) **Culture:** on blood agar, typical colonies appear in 18 hrs at 37 °C after several days we could see the haemolysis and pigment production. In case of mixed flora, we could use media containing 7.5% NaCl (Staph 110 and mannitol salt agar. On the latter they develop yellow color). On Baird Parker agar colonies appear black surrounded by clear halo.
- d) **Coagulase test:** it separates staphylococci into two groups; coagulase positive and coagulase negative.
- e) **Catalase test:** it differentiates the staphylococci, which are positive, from the streptococci, which are negative.
- f) **Serological tests:** Antibodies to teichoic acid specially can be detected in prolonged, deep infections (e.g. endocarditis). These serologic tests have low practical value.
- g) **PCR.**

❖ Epidemiological (nosocomial) infections

In case of hospital acquired infection or an epidemic spread out in the hospital, due to *S. aureus* it is so important to diagnose the type or strain of *S. aureus* that cause the infection. This goal can be achieved by following a kind of tests known as "EPIDEMIOLOGICAL MARKERS":

- 1) **Biotyping**: not all isolates that had been isolated from different sites, are 100% identical in all characters.
- 2) **Phage typing**: there are about 100 phages (can infect more than 22 specified types of *S. aureus*) classified into four groups called lytic groups.
- 3) **Antibiogram**: similar or identical patterns of resistance to antibiotics can also be used as presumptive evidence of the relatedness of strains in epidemiological tracing. Methicillin resistance is very important.
- 4) **Serotyping**: there are more than 30 specific antigens (agglutinogens)
- 5) **Genetic methods**: DNA hybridization or plasmid profile.
- 6) **Polymerase chain reaction (PCR)**

❖ Treatment

Gram positive bacteria usually treated by penicillin but that were until 1970 then resistant strains appeared and penicillin becomes to cause allergy. So we can use: Cephalosporin, Methicillin, Oxacillin or an aminoglycoside. Refampin (in case of endocarditis, since it can reach inside the valve). In case of methicillin resistance vancomycin is appropriate alternative.

❖ Sources of infections

1. **Patients** with lesions are discharging staphylococci.
2. **Healthy carriers**: they are healthy, but carrying the microorganisms, hence they considered as source of infection, especially food handler.

3. **Animals:** some of them have strains that affect humans and could be transmitted to human by direct contact with animals raised at homes and cause disease.

❖ **Prophylaxis**


No vaccine had been developed against *S. aureus* because antigens are very wide, toxins have variable mode of action. However, cleaning is very important to prevent infection.

II- COAGULASE NEGATIVE STAPHYLOCOCCI (CONS)

Interest had increased in CONS when we started using clinical prosthetic devices. They may cause wound infection, UTI, bacteraemia, catheters infections, vascular grafts, and infections of prosthetic devices.

☒ *S. epidermidis*: It a major skin inhabitant it causes endocarditis, colonization of prosthetics, bacteraemia, wound infections, and UTI especially in elderly hospitalized patients, sensitive to novobiocin.

☒ *S. saprophytics*: Causes 10-20% of primary Urinary Tract Infections (UTI) in young women (16-35 yrs). They are resistant to novobiocin.



Lecture 9: streptococci

Dr. Harith Jabbar Fahad
DEPARTMENT OF BIOLOGY

Gram positive cocci**Genus: *Streptococcus*****▪ General characteristics**

Gram positive cocci that appear in chains or clusters, most species are facultative anaerobes except one group "strict aerobes" which inhabits the intestinal tract and female genital tract. All streptococci are catalase negative. Some streptococci elaborate capsular polysaccharide. Hair-like pili project through the capsule of group A streptococci. The pili consist partly of M protein and are covered with lipoteichoic acid.

They are fastidious requiring many amino acids, vitamins, purine and pyrimidine bases. Primary habitat of upper respiratory tract of human.

▪ Classification

1 - Depending on haemolytic behaviour on blood agar:

a. Alpha haemolytic / incomplete haemolysis / viridans group.

b- Beta haemolytic / complete haemolysis / major human pathogens.

c- Gamma haemolytic / no haemolysis / not primary pathogens.

2- Rebecca Lancefield and co-workers / classification depending on antigenic characteristics of cell wall; carbohydrate C- substance in **beta haemolytic** streptococci / serological groups **A-V** (except I and J).

3- Biochemical reactions and resistant to physical and chemical factors.

4- Molecular genetics and ecologic features.

▪ Group A streptococci (*Streptococcus pyogenes*)**❖ Antigenic structure**

- 1- **Capsule:** most group A, B, and C strains produce capsules composed of hyaluronic acid. Those who has capsule can impede the phagocytosis. This capsule is antigenic but not immunogenic.
- 2- **Cell wall proteins**
 - **M protein:** this substance is a major virulence factor of group A *S. pyogenes*. When M protein is present, the streptococci are virulent, and they are able to resist phagocytosis. *The M protein (found in fimbriae) binds fibrinogen from serum and blocks the binding of complement to the underlying peptidoglycan.* It is immunogenic. There are more than 80 different types of M protein. Strains of *S. pyogenes* that produce certain M protein types are rheumatogenic, whereas strains of *S. pyogenes* that produce other M protein types are nephritogenic.
 - **T protein:** this antigen has no relationship with virulence of Streptococci. It permits differentiation of certain types of streptococci, especially those whom isolated from impetigo patients, by agglutination with specific antisera.
 - **Cell surface proteins: R protein** is used as an epidemiologic marker and has no known role in virulence. **F protein** (fibronectin binding), **G protein** binds the Fc portion of antibodies, and **P substance**, which probably make up most of the streptococcal cell body.
- 3- **Cell wall carbohydrates:** according to them, Streptococci are classified into serological groups (Lancefield groups). The serologic specificity is determined by an amino sugar. For group A Streptococci, this is rhamnose-*N*-acetylglucosamine; for group B, rhamnose-glucosamine polysaccharide; for group C, rhamnose-*N*-acetylgalactosamine; for group D, glycerol teichoic acid containing D-alanine and glucose; for group F, glucopyranosyl-*N*-acetylgalactosamine. **Group A sugar (also called**

C-substance) can cross-react with heart valves glycoprotein.

C-substance found to cause arthritis to some animals.

- 4- **Cell wall peptidoglycan:** causes fever, lysis of red blood corpuscles in rabbits and other animals, and dermal necrosis.
- 5- **Cytoplasmic membrane antigens:** cross- reacts with tissues of heart, kidneys, and connective tissues. Group D specific antigen belongs to the cytoplasmic membrane antigens.

▪ **Extracellular streptococcal products**

The Streptococci elaborate extracellular products in vitro. Many of these products are believed to be important in the pathogenesis of streptococcal infections. They are used in diagnosing such infections.

1- Pyrogenic exotoxin (formerly known as Erythrogenic toxin): Three **streptococcal pyrogenic exotoxins (SPE)**, are recognized; A, B and C. In the non-immune individual, it causes rash that is characteristic of scarlet fever.

2- Cardiohepatic toxin and nephrotoxin: a low molecular weight compound excreted by virulent strains of Streptococci is capable of producing lesions in the heart and liver tissue when injected into susceptible animals.

3- Haemolysins: the haemolytic activity of many streptococcal strains is due to the production of two distinct extracellular haemolysins called streptolysin O (SLO) and streptolysin S (SLS).

☒ **SLO:** is oxygen labile and immunogenic. Detection of antistreptolysin O antibodies is extensively used in the diagnosis of streptococcal infections. SLO has been implicated as a factor in the pathogenesis of rheumatic fever, a post streptococcal complication.

☒ **SLS:** is oxygen stable, responsible for the zone of beta haemolysis seen around surface colonies of streptococci on blood agar plates. It is capable of lysing mammalian red blood corpuscles and leukocytes, it is non-immunogenic. Intravenous injections of SLS in rabbits cause intravascular haemolysis and acute liver necrosis.

3- Spreading factors:

a. Hyaluronidase: can digest host connective tissue hyaluronic acid, as well as the organism's own capsule. Depolymerization could therefore enhance the spread of streptococci in the tissue. However, it could be used as treatment in certain cases.

b. Protease: a proteolytic activity has been shown in strains causing soft tissue necrosis or toxic shock syndrome.

c. Streptokinase:

produced by most of group A streptococci. It can interact with the proenzyme plasminogen of human serum, converting it to plasmin. Plasmin can digest fibrin and other serum factors important in the formation of blood clots. This activity is now used in the treatment of acute myocardial infarction.

d. Nucleases (streptodornases A-D): At the site of infection, the host's inflammatory response results in the accumulation of nuclear exudates from dead or injured white blood cell. Virulent streptococci can produce nucleases such as ribonuclease and deoxyribonuclease that will digest DNA and RNA of the exudates thus facilitating the spreading of streptococci.

e. Other extracellular products: such as C5a peptidase: is an extracellular enzyme that degrades complement component C5a, the main factor that attracts phagocytes to sites of complement deposition.

This large repertoire of products is important in the pathogenesis of *S. pyogenes* infections. Even so, antibodies to these products are relatively insignificant in protection of the host.

▪ Pathogenesis

Ninety percent of streptococcal diseases caused by group A beta haemolytic Streptococci. Streptococcal diseases are divided into two categories:

I- Suppurative diseases:

- a. **Impetigo**: a highly contagious skin disease found primarily in children, involves the infection of *epidermal layers of skin*. The infection begins as small blisters that can spread to adjacent areas.
- b. **Cellulites**: inflammatory condition associated with streptococcal invasion of *subcutaneous tissue*. This type of disease results in gangrene and invasion of blood stream.
- c. **Erysipelas**: involves the infection of *dermis* characterized by a spreading inflammation with massive brawny oedema and a rapidly advancing margin of infection.
- d. **Necrotizing fasciitis (streptococcal gangrene)**: this is an infection of the subcutaneous tissues and fascia. There is extensive and very rapidly spreading necrosis of the skin and subcutaneous tissues. Group A streptococci that cause necrotizing fasciitis have sometimes been termed "**flesh-eating bacteria**".
- e. **Puerperal fever**: uterine infection that frequently accompanies delivery when aseptic techniques are not followed.

f. Streptococcal sore throat: in children and adults, the disease is acute and is characterized by intense nasopharyngitis, tonsillitis, and intense redness and oedema of the mucous membranes, with purulent exudates; enlarged, tender cervical lymph nodes; and (usually) high fever.

g. Sepsis: infection of traumatic or surgical wounds with streptococci results in sepsis or surgical scarlet fever.

h- Toxic shock: is caused by a few strains that produce a toxic shock-like toxin.

II- Non suppurative diseases

a. Scarlet fever: associated with the formation of erythrogenic toxin by group A streptococci.

b. Rheumatic fever: M protein cross reacts with sarcolemma. Antibodies cross-react with heart tissue, fix complement, and cause damage accompanied by inflammation of the joints. Highest incidence found in age group 5-19 years.

c. Acute glomerulonephritis (AGN): Antigen-antibody complexes may be deposited in kidney, fix complement, and damage glomeruli. Only a few M-types are nephritogenic.

d. Erythema nodosum: skin condition, small red nodules appear under the surface of the skin. It could be the result of hypersensitivity to the peptidoglycan of the streptococcal cell wall.

▪ Identification

Preliminary identification:

1 - Haemolytic behaviour. Cultures on sheep blood agar plates are the gold standard.

- 2- Bacitracin sensitivity: group A streptococci are sensitive to Bacitracin, while group B streptococci are resistant.
- 3- Fluorescent antibody.
- 4- Phadebact coagglutination test: Major beta haemolytic groups A, B, C, and G detected by this kit.
- 5- Streptex: latex particles are conjugated to group specific streptococcal antibodies mixed with unknown growth of streptococci result in agglutination.
- 6- Directly from throat swab (large numbers of streptococci required) is mixed with nitrous acid (extraction) in combination with coagglutination. The test required 30 minutes.
- 7- PCR.

▪ **Diagnostic test for post streptococcal diseases:**

- 1- Determination of anti streptolysin O titer (ASOT). An elevated ASO is a useful diagnostic marker for rheumatic fever. In contrast to rheumatic fever, antistreptolysin O (ASO) titers are low in acute glomerulonephritis.
- 2- Streptozyme test (slide agglutination test): Sheep red blood corpuscles are sensitized with extracellular products from group A streptococci mixed with diluted patient serum results in agglutination. The extracellular products are Hyaluronidase (HA), deoxyribonuclease (DNase), nicotinamide adenine dinucleotidase (NADase), and streptokinase (SK).

▪ **Treatment and prevention**

Penicillin is the drug of choice for treatment of group A beta haemolytic streptococci. In case of individuals sensitive to **penicillin**, **erythromycin** is a suitable alternative. **Vaccines** are prepared from streptococcal type specific M protein against group A infections.

- **Other groups**

1- **Group B Streptococci:** they found in oral cavity, intestinal tract and vagina *S. agalactiae*, which surrounded by a polysaccharide capsule. They cause neonatal meningitis by transporting to the baby during delivery, UTI, and puerperal fever.

2- **Group C Streptococci:** cause infections in many animals' species but rarely in human. *S. dysgalactiae subsp. equisimilis* cause human illnesses such as endocarditis, pneumonia, meningitis, and wound infections. Outbreaks of disease associated with unpasteurized milk or cheese made from unpasteurized milk. *S. equi* causes strangles in horses. *S. equi subsp. zooepidemicus* infections are infrequent in human e.g. meningitis and pneumonia.

3- **Group D Streptococci:** alpha or gamma haemolytic on sheep blood agar, of intestinal origin (Enterococci): *E. faecium*, *S. bovis*, *S. equinus*, *Enterococcus durans*, *Enterococcus faecalis*, the latter is of medically important since it causes UTI and abdominal lesions.

- **Non-Lancefield Group Streptococci**

Includes **the viridans Streptococci** (*S. mutans*, *S. sanguis*; *S. salivarius*; *S. mitis*) and *S. pneumoniae*.

Streptococcus pneumoniae

Formerly, *Diplococcus*. Inhabitants of the upper respiratory tract, gram positive cocci that occur singly, in pairs, or in chains. The clinical specimens are lancet shaped and surrounded by a capsule, only encapsulated (smooth form) strains are virulent. On repeated subculture in the laboratory, the capsule is lost (rough form). The pneumococci are facultative anaerobes that

are very fastidious in their cultural requirements. Some strains need and elevated level of CO₂ (5-10 %) for initial isolation.

The addition of blood to culture media supplies the enzyme catalase. Under aerobic conditions pneumococci produce hydrogen peroxide, which can be toxic to the pneumococci. The enzyme catalase acts to remove the accumulated hydrogen peroxide.

On blood agar pneumococcal colonies **exhibit alpha haemolysis** and closely resemble colonies of alpha haemolytic Streptococci.

☒ **Virulence factors**

1- Capsular polysaccharide: the primary virulence factor of the pneumococci. The capsule prevents binding of antibody to the cell wall of the pneumococcus and thus inhibits phagocytosis.

2- Pneumolysin: all pneumococci produce pneumolysin. This protein is related to *S. pyogenes* streptolysin O. Pneumolysin is toxic to pulmonary endothelial cells. Pneumolysin deficient pneumococci are less virulent than the toxin producing isolates. Most patients with pneumococcal disease exhibit an antibody response to pneumolysin and pneumolysin immunization is protective in animals.

3- Pneumococcal surface protein A (PspA): found on all pneumococci, and highly variable both immunologically and in molecular mass. Passive immunization with anti PspA is protective.

4- Neuraminidase: like pneumolysin, is released upon autolysis. Despite the involvement of this enzyme in the colonization by other bacteria, there is no proven role in pneumococcal virulence.

5- SIgA protease: all pneumococci produce SIgA protease, which is a property shared with other species causing pneumonia and meningitis. This enzyme could play a part in establishment of the microorganisms in the nasopharynx.

☒ Infections produced by Pneumococci

1- Primary infections: caused by certain serotypes and responsible of approximately 75 % of pneumonia cases and responsible of more than half of lethal cases caused by pneumococcal bacteraemia. Lobar pneumonia, Septicaemia, Peritonitis, Purulent meningitis and purulent otitis are examples of primary infections.

2- Secondary infections

Produced by any serotype, 40-70% of the population is carriers of pathogenic *S. pneumoniae*. Many authorities believe that pneumonia and related pneumococcal infections are acquired **endogenously** through **lowered host resistance** rather than exogenously by direct contact.

☒ Laboratory diagnosis of pneumococci

Specimens: sputum, laryngeal swabs, transtracheal aspirates, blood, and CSF. A definitive diagnosis of pneumococcal pneumonia can be made only if the microorganisms are isolated directly from the blood or from other clinical specimens by plating onto blood agar. Alternatively, by identification of specific pneumococcal antibodies by counter **Immunelectrophoresis** of body fluids: pleural fluid, blood, and urine.

The pneumococci can be differentiated from other alpha haemolytic streptococci by the following procedures:

1- Quellung reaction: it is the most accurate and specific test of identification of the pneumococci. Sputum or exudative material is spread on a slide and mixed with antiserum against type specific polysaccharide or the polyvalent antiserum results capsular swelling. A capsular halo can be observed around the diplococci.

2- Optochin susceptibility test: optochin is an antimicrobial drug derived from quinine. When disks containing the drug are placed on blood agar

previously seeded with pneumococci, zones of inhibition can be observed around the optochin. Alpha haemolytic Streptococci are resistant to optochin.

3- Bile solubility test: the pneumococci produce an enzyme (amidase) that cleaves specific covalent bonds in the peptidoglycan layer. This enzyme is activated by bile or bile salts such as sodium deoxycholate solution are added to a broth culture of pneumococci, the cells are rapidly lysed. While, Alpha haemolytic streptococci are not lysed by bile.


4- Mouse virulence test: most pneumococcal strains, when injected intraperitoneally into mice, can induce death of animals within 24-48 hours. Alpha haemolytic streptococci, used under the same procedures, are not lethal to mice.

☒ Treatment of pneumococci

Penicillin is the drug of choice. In **penicillin** allergic patients, **erythromycin** or one of its derivatives **azithromycin**, can be used. **Vancomycin** is the drug of choice for the **penicillin** resistant pneumococci.

☒ Prevention of pneumococci

In **1983**, a polyvalent vaccine containing polysaccharide antigen from 23 types of *S. pneumoniae* was recommended for children over 2 years. In **1998**, a vaccine containing pneumococcal polysaccharide **coupled to a carrier protein (diphtheria toxoid)** as the immunogen was shown to be effective in young children. The vaccine contained the polysaccharide of the seven most common pneumococcal serotypes. This vaccine was approved by the FDA in **2000**.



GRAM NEGATIVE COCCI

Lecture 10



THE GRAM NEGATIVE COCCI and related organisms

I- *Neisseria*

General characteristics:

1. All are fastidious Gram negative diplococci with a bean shaped configuration, the flat or concave sides are adjacent.
2. Flagella and swimming motility are absent. The cells are nonmotile in liquid media but surface-bound motility (“twitching motility”) is frequently observed.
3. All species are aerobic or facultatively anaerobic. The pathogens grow better in 5 to 10% CO₂.
4. All species produce cytochrome oxidase (oxidase positive).

The medically important members are *N. meningitidis* (**the meningococcus**) and *N. gonorrhoeae* (**the gonococcus**), both are pathogenic for humans (obligate parasite) and typically are found associated **with or inside polymorphonuclear cells**. The organisms are sensitive to low temperature and drying.

Neisseria meningitidis (the meningococcus)

Isolated by Weichselbaum in 1887

❖ Antigenic composition of meningococci

Antigens are organized in three ways:

1. **Serogroups:** meningococci are divided into at least 12 serogroups based on **chemical and structural characteristics of capsular polysaccharides**. Serogroups: A, B, C, X, Y, Z, 29E, and W-135 are of medical important. A, B, and C produce epidemics. Serogroups X, Y, and Z produce meningitis, single cases; and found in carriers of the disease.
2. **Serotypes:** serotyping based on **outer membrane proteins of the cell wall (e.g. porin proteins)**. These serotypes have become

important in studies of the epidemiology of infection and in the development of new vaccines.

3. **Immunotypes:** immunotyping based on **lipopolysaccharides**. Neisserial LPS is distinguished from enteric LPS by its highly-branched basal oligosaccharide structure and the absence of repeating O-antigen subunits. For these reasons, neisserial LPS is referred to as **Lipooligosaccharides (LOS)**.

❖ Pathogenicity of meningococci

The most important virulence factors contributing to disseminated disease are pili, IgA1 protease, LOS, outer membrane proteins, and capsule polysaccharides.

Infection with *N. meningitidis* has two presentations, **meningococemia**, characterized by skin lesions, and **acute bacterial meningitis**. The fulminant form of disease (with or without meningitis) is characterized by multisystem involvement and high mortality.

Infection is by aspiration of infective bacteria, which attach to epithelial cells of the nasopharyngeal and oropharyngeal mucosa by the aid of pili, cross the mucosal barrier, and enter the bloodstream. It is not clear whether blood-borne bacteria may enter the central nervous system and cause meningitis.

The mildest form of disease is a transient bacteremic illness characterized by a fever and malaise; symptoms resolve spontaneously in 1 to 2 days. The most serious form is the fulminant form of disease complicated by meningitis.

Chills, fever, malaise, and headache are the usual manifestations of infection. Signs of meningeal inflammation are also present.

The majority of people who contract bacterial meningitis and meningococcal septicemia survives and make a full recovery; however,

some are left with major residual effects such as deafness, mental retardation, and behavioral defects.

❖ **Diagnostic laboratory tests of meningococci**

- ☒ **Specimens:** specimens of blood are taken for culture, and specimens of spinal fluid (lumbar puncture) are taken for smear, culture, and chemical determination.
- ☒ **Smears:** gram stained smears of the sediment of centrifuged spinal fluid often show typical *Neisseriae* **within** polymorphonuclear leukocytes or extracellularly.
- ☒ **Culture:** specimens are cultured on media such as chocolate agar, Mueller-Hinton agar or modified Thayer-Martin medium with antibiotics (vancomycin, colistin, amphotericin). Yielded cultures can be further identified by carbohydrate fermentation reactions.
- ☒ **Oxidase test.**
- ☒ **Sugar fermentation test.**
- ☒ **Serology:** it usually performed on CSF, such as capsular swelling, fluorescent antibody staining, and counter Immunoelectrophoresis (CIE). Negative capsular swelling tests and negative CIE test results ***do not necessarily rule out*** a meningococcal infection because serogroups other than A and C frequently possess insufficient capsular antigens to be detected by these methods.

❖ **Prevention and control of meningococci**

A) Chemotherapy

Antimicrobial drugs; Sulpha drugs were drug of choice until bacteria developed resistance against these drugs. Therefore penicillin is the drug of choice. In cases of penicillin allergic patients, chloramphenicol or ampicillin with moxalactam could be used.

Ciprofloxacin, ceftriaxone or rifampicin may be used prophylactically as a means of preventing the disease state in carriers.

B) Active immunization

Immunizing materials have been developed against serogroups A, C, Y, and W-135 meningococci. However, Group C vaccine does not induce protective levels of antibody in children younger than 2 years. Group B are poorly immunogenic.

❖ Epidemiology

Meningococcal are endemic worldwide. They inhabit mainly the nasopharynx and transmitted via droplets or close kissing contact. Of the young adult population, 10-30% carry meningococci. Carriage is highest among enclosed populations. The most vulnerable age groups are children aged < 5 years, particularly those < 2 years, followed by teenagers and young adults. This disease is more common during autumn and winter seasons.

***Neisseria gonorrhoeae* (gonococcus GC)**

Isolated in 1879 by Neisser. They considered as one of the most common sexually transmitted pathogen.

❖ Antigenic structure of gonococci

Neisseria gonorrhoeae is antigenically heterogeneous and capable of changing its surface structures in vitro, and presumably in vivo, to avoid host defenses. Surface structures include: pili, porins proteins, opacity associated proteins (opa), LOS and others. Gonococci do not have capsular polysaccharides.

❖ Virulence factors of gonococci

The following factors are currently regarded as possible agents of gonococcal pathogenicity:

1) Pili: mediate initial attachment of gonococci to epithelial cells.

Gonococci produce four colony forms: T1, T2, T3, and T4. Only the

bacteria in T1 and T2 colonies are piliated, and are regarded as virulent because they produce infection in volunteers.

- 2) **Por proteins (protein I):** when the gonococcal membrane is in intimate contact with the host cell membrane, the Por protein is transferred to the host cell, resulting in alterations in ionic permeability of the host cell plasma membrane. May contribute to the intracellular survival of gonococci inside of neutrophils.
- 3) **Opa (protein II):** afimbrial adhesins that mediate firm attachment of gonococci to epithelial cells.
- 4) **IgA protease:** cleaves and inactivates the SIgA subclass IgA1. Nonpathogenic *Neisseria* species do not produce IgA protease.
- 5) **Epithelial endocytosis:** the gonococci attach to the epithelial cells of the cervix, are subjected to endocytosis, multiply within the cell, and are protected from the phagocytic activities of leukocytes.
- 6) **Endotoxin (LOS):** mediates most of the toxic damage in the epithelial cells, and regulates complement activation on the surface of the organism. It also is implicated in the attachment of gonococci to host cells by piliated and nonpiliated organisms. The sialylation of the LPS results in the conversion of a serum-sensitive organism to serum-resistant. Gonococcal endotoxin is probably responsible for some disseminated gonococcal infections (DGI).

❖ **Gonococcal diseases**

A. Gonorrhoea

Gonorrhoea is one of the most common bacterial venereal diseases. Complicating the high incidence of disease is the increasing appearance of multiple antibiotic resistance. The disease is generally spread via sexual activity.

Disease in men occurs after an incubation period of from 2-14 days. Onset of disease is usually marked by mild discomfort in the

urethra, followed a few hours later by dysuria and a purulent yellowish green urethral discharge. In women, symptoms usually begin within 7-21 days after infection. A significant portion of infected women is asymptomatic (70 - 80%). However, onset is sometimes severe, with dysuria and vaginal discharge. The cervix and deeper reproductive organs are the sites most frequently infected.

Complications, in men, postgonococcal urethritis, a common sequela, actually results from infection with other organisms (e.g., *Chlamydia trachomatis*). Epididymitis is uncommon and usually unilateral. Prostatitis and urethral stricture are less common. Ascending infection may result in infertility. In women, salpingitis (pelvic inflammatory disease) is the most important clinical problem. This syndrome has two important consequences: 1) sterility and ectopic pregnancy; and 2) susceptibility to chronic infections. DGI with bacteremia is more common among women than men.

B. Extragenital infections

- i. **Pharyngitis and conjunctivitis:** Infants born to a mother with cervicovaginal gonorrhea may develop gonococcal conjunctivitis.
- ii. **Gonococcal arthritis** may be preceded by symptomatic bacteremia. The onset typically is acute, with fever, severe pain and limitation of movement in one or a few joints.

❖ Pathogenesis of gonococci

Gonococci initially attach to host epithelial cells via **pili**. Closer attachment is then quickly mediated by the **Opa** protein. Secretory IgA protease protects the organisms from antibodies present at the mucosal surface. Some epithelial cells are damaged by gonococcal LOS, but **others** are probably **invaded** by the organism. After attachment and initial colonization, the sequence of events probably includes: **1)** Entry of gonococci

into the host cell by endocytosis. **2)** Intracellular replication inside the endocytic vesicle. (Host cell killing of bacteria within the vesicles is inhibited by the membrane perturbing activities of **Por**). **3)** Transport of the vesicle to the basal of the cell, fusion with the cell membrane, and release of the gonococci into the subepithelial tissue, the lamina propria. **4)** Multiplication in the lamina propria aided by iron acquisition systems. **5)** The gonococci have opportunities to spread because of the proximity of the lamina propria to regional lymphatics and blood vessels; DGI happens in approx. 1% of cases.

❖ Laboratory diagnosis of gonococci

Laboratory identification procedures include the following:

- 1) Specimens:** The specimens selected for diagnosing gonorrhea depend on the gender, age and sexual preference of the patient. Urethral and cervical specimens routinely are collected from men and women, respectively. However, the organism may be isolated from blood, the nasopharynx, skin lesions, CSF and the anal canal of homosexual.
- 2) Direct examination:** Gram stain of purulent materials may reveal Gram-negative diplococci in polymorphonuclear leukocytes.
- 3) Culture:** Specimens are cultured on media such as supplemented chocolate agar or Thayer-Martin medium in an elevated CO₂ environment". Blood cultures and cultures of synovial aspirates or skin lesions should be attempted for patients with suspected disseminated gonococcal infection (DGI).
- 4) Oxidase test.**
- 5) Superoxol test:** catalase test using 30% H₂O₂ was used to differentiate *N. gonorrhoeae* from other *Neisseria* species.
- 6) Carbohydrate fermentation.**
- 7) Serological tests:**
 - a. Direct fluorescent antibody technique.

b. Staphylococcal coagglutination technique.

8) Molecular methods: The microbiological diagnosis of gonorrhea based on culture on selective medium produces about 80 - 95% sensitivity, with false-negative results attributed to poor specimen storage, transport problems, and inhibition of growth by the components of selective media. As an alternative diagnostic test, molecular techniques have been developed and offer the promise of eliminating transport and specimen collection issues, which are believed to affect test sensitivity in the field setting.

❖ **Treatment of gonococci**

The recommended treatment for uncomplicated infections is a third-generation cephalosporin or a fluoroquinolone plus an antibiotic (e.g. doxycycline or erythromycin) effective against possible coinfection with *Chlamydia trachomatis*. Sex partners should be referred and treated. The current **CDC Treatment Guidelines** recommend treatment of all gonococcal infections with antibiotic regimens effective against resistant strains. The recommended antimicrobial agents are ceftriaxone, cefixime, ciprofloxacin, or ofloxacin.

Prophylaxis of "ophthalmia neonatorum" can be achieved by local application of 0.5 % erythromycin ophthalmic ointment or 1% tetracycline ointment.

No vaccine was prepared until now. The patients failed to develop a solid immunity.

II- *Acinetobacter*

Includes many species of gram negative diplococci or coccobacilli. Widely distributed in nature and are part of the normal human flora (skin).

Acinetobacter calcoaceticus an opportunist causes variety of infections especially hospital associated infections; Wound infections, Meningitis, Pneumonia, Bacteraemia, and Urethritis.

III- *Moraxella*

The moraxella group includes six species. They are nonmotile, nonfermentative, and oxidase positive. On staining, they appear as small gram negative bacilli, coccobacilli, or cocci. They are members of the normal flora of the upper respiratory tract and occasionally cause bacteraemia, endocarditis, conjunctivitis, meningitis, or other infections. Most of them are susceptible to penicillin.

A) *Moraxella catarrhalis*

It was previously named *Branhamella catarrhalis* and before that *Neisseria catarrhalis*. It is a member of the normal flora in nasopharynx. It causes bronchitis, pneumonia, sinusitis, otitis media, and conjunctivitis.

B) *Moraxella lacunata*: Causes conjunctivitis.